

## Cryopreservation of *in situ* cool stored buffalo (*Bubalus bubalis*) epididymal sperm

Barati, F.<sup>1\*</sup>; Khaksary Mahabady, M.<sup>2</sup> and Mohammadi, Gh.<sup>1</sup>

<sup>1</sup>Department of Clinical Sciences, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran; <sup>2</sup>Department of Basic Sciences, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran

\*Correspondence: F. Barati, Department of Clinical Sciences, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran. E-mail: baratif@scu.ac.ir

(Received 9 Mar 2009; revised version 20 Jun 2009; accepted 4 Jul 2009)

### Summary

The objectives of this study were to investigate the effects of testis-epididymis cool storage on viability and progressive motility of buffalo epididymal sperm (EP) and to compare the influences of two basic semen extenders on post thaw EP viability and progressive motility. Abattoir collected buffalo testicles were allotted to three storage times (0 h: n = 10; 24 h: n = 10 and 48 h: n = 12). Following storage, isolated sperm were subjected to cryopreservation with two different cryoprotective media: whole cow milk-7% glycerol (MG) or egg yolk-tris-citrate-7% glycerol (EYG). Pre freeze and post thaw sperm progressive motility and viability were evaluated. Results indicated that viability and progressive motility of sperm decreased after 24 h cool storage of the epididymis ( $P < 0.05$ ). There was no difference between 24 and 48 h of storage on sperm viability ( $P > 0.05$ ), but progressive motility decreased across storage times (24 h *versus* 48 h:  $P < 0.05$ ). Cryopreservation severely influenced viability and progressive motility of EP ( $P < 0.05$ ). Milk-7% glycerol protected viability and progressive motility of EP against cold shock more efficiently than EYG ( $P < 0.05$ ). The results of this study demonstrate that it is possible to preserve buffalo EP within the epididymis at 4°C short term, but that it has poor freezability upon recovery by basic semen freezing protocols.

**Key words:** *Bubalus bubalis*, River Buffalo, Epididymal sperm, Cryopreservation, Cool storage

### Introduction

There is a considerable population (>500,000 head) of river buffalo (*Bubalus bubalis*) in Iran which play a significant role in rural life by producing milk and meat while tolerating the impact of harsh environmental conditions (Naderfard and Qanemy, 1997). This species is not in danger of extinction in Iran, but the rate of fertility, which is directly correlated with production, is low (Barile, 2005). Although buffalo semen has been stored successfully in liquid and frozen states (Sansone *et al.*, 2000), artificial insemination (AI) has not yet been developed mainly due to oestrus detection issues (Barile, 2005). With the new approaches developed for the management of ovulation (e.g., Ovsynch program), assisted reproductive technologies (ART) could help to overcome reproductive insufficiency in the buffalo species (Barile, 2005).

Epididymal sperm (EP) have been successfully used for AI and *in vitro* production of embryos (IVP) in several species (Kikuchi *et al.*, 1998; Hori *et al.*, 2004; Hori *et al.*, 2005). Motile sperm have been isolated from cool stored epididymis of boar (Kikuchi *et al.*, 1998), cattle (Nichi *et al.*, 2007), dog (Stilley *et al.*, 2000; Yu and Leibo, 2002), stallion (Braun *et al.*, 1994; Bruemmer *et al.*, 2002), African buffalo (*Syncerus caffer*) (Bartels *et al.*, 1999), goat (Blash *et al.*, 2000), ram (Kaabi *et al.*, 2003), camel (Wani *et al.*, 2005), Iberian red deer (*Cervus elaphus hispanicus*) (Soler *et al.*, 2005; Fernández-Santos *et al.*, 2006) and Sika deer (*Cervus nippon*) (Hishinuma *et al.*, 2003). Tomcat EP (Hermansson and Axner, 2007) has been diluted in a chemical medium and was motile up to 72 h under refrigerator condition.

Influences of different types (Herold *et al.*, 2004; Wani *et al.*, 2005) and concentrations (Herold *et al.*, 2004;

Fernández-Santos *et al.*, 2006; Herold *et al.*, 2006; Morton *et al.*, 2007) of cryoprotectants, additives to freezing medium (Herold *et al.*, 2004) and determination of equilibration time (Herold *et al.*, 2004; Herold *et al.*, 2006) have been investigated for EP cryopreservation. Morton *et al.* (2007) compared citrate based, tris based and lactose based diluents for alpaca EP freezing. Herold *et al.* (2004, 2006) used AndroMed® (a chemical defined medium) and Triladyl™ (an egg yolk based medium) for cryopreservation of African buffalo (*Syncerus caffer*) EP. They reported the optimal equilibration time of 4 h for African buffalo (*Syncerus caffer*) EP freezing (Herold *et al.*, 2006). Cool stored epididymal Iberian red deer (*Cervus elaphus hispanicus*) (Fernández-Santos *et al.*, 2006) and stallion (Braun *et al.*, 1994; Bruemmer *et al.*, 2002) sperm were successfully cryopreserved.

This study was undertaken to determine 1) the effect of duration of cool storage of excised epididymes on viability and progressive motility of river buffalo EP, and 2) the influence of two basic diluents (milk-7% glycerol (MG) and egg yolk-tris-citrate-7% glycerol (EYG)) on post thaw viability and progressive motility of buffalo EP.

## Materials and Methods

Tyrodes lactate (TL) solution components, fructose, citric acid, and glycerol were purchased from Merck, Germany. *N*-2-hydroxyethylpiperazine-*N*-2-ethane-sulfonic acid (HEPES) buffer, bovine serum albumin (BSA; Fraction-V), pyruvic acid, Tris (TRIZMA®BASE), phenol red and gentamycin were provided from Sigma-Aldrich, St. Louis, USA. Other materials are specified through the text.

### Media preparation

Sperm TL medium was tyrodes lactate medium (100 mM NaCl, 3.1 mM KCl, 25 mM NaHCO<sub>3</sub>, 0.29 mM NaH<sub>2</sub>PO<sub>4</sub> H<sub>2</sub>O, 21.6 mM Na Lactate, 2.1 mM CaCl<sub>2</sub> 2H<sub>2</sub>O, 0.4 mM MgCl<sub>2</sub> 6H<sub>2</sub>O and 10 mM HEPES buffer; Parrish *et al.*, 1988) plus 0.006 g/ml BSA, 1 mM sodium pyruvate, 25 µg/ml gentamycin and 10 mg/L phenol red (pH =

7.4 and osmolarity = 290-305 mOsm/L). The cryoprotective media (pH = 6.8-7.4) were egg yolk-citrate-glycerol (199.8 mM TRIZMA®BASE, 45 mM citric acid, 55.5 mM D-fructose, 20% (v/v) egg yolk 7% (v/v) glycerol and 0.5 mg/ml gentamycin; Salisbury and Fuller, 1941) and cow milk (Choopan dairy industries, Tehran, Iran) - glycerol (whole cow milk with 7% (v/v) glycerol and 0.25 mg/ml gentamycin; Zaugg and Almquist, 1972). Each of the media was prepared separately in two parts (part A: media without glycerol and part B: media with 14% glycerol). One batch of each part was prepared and frozen for subsequent use in 10 ml aliquots. On the day of use, the parts A were thawed and warmed to room temperature (22-25°C) and the parts B were thawed at 4°C until liquefied.

### Testes preparation and EP isolation

Mature buffalo bull (>2.5-year-old) testes were collected from the local abattoir, placed on ice and transported to the laboratory. The testes were separated from each other and allotted to three different times of cool storage (0 h: n = 10; 24 h: n = 10; 48 h: n = 12). After storage, each testicle was brought to room temperature in less than 30 min and cauda epididymis was dissected from testis-epididymis, minced and placed into a 60 mm tissue culture dish containing sperm TL medium for 15 min. Then the tissue was removed and the remaining medium was centrifuged (Behdad®, Tehran, Iran) at 700 g for 6 min to create a sperm pellet. Supernatant was removed and the isolated sperm analyzed for sperm concentration, viability and progressive motility. Finally, a sperm suspension (300 × 10<sup>6</sup> sperm/ml) was created by adding sperm TL medium. The diluted EP (provided from each testis-epididymis) was subjected to cryopreservation.

### Cryopreservation and thawing procedures

The two step dilution procedure was used for sperm cryopreservation (Salisbury and Fuller, 1941). The sperm suspension was divided equally and added to part A of the diluents (EYG and MG) at room

temperature to create a  $30 \times 10^6$  sperm/ml sperm dilution. The extended sperm was placed in a 25°C water jacketed tube and cooled to 4°C within 2.5 h in refrigerator. An equal volume of part B of the diluents (EYG and MG) was added in two steps (1:2 and 1:2 of original volume) with a 15 min interval, resulting in a final concentration of 7% glycerol. The glycerolized samples were equilibrated for an additional 1 h at 4°C before being loaded into straws (0.25 ml french straws; IMV, France). The filled straws were sealed, placed 2-4 cm above the liquid nitrogen (LN2) level for 8 min, and then immersed into LN2. At least 48 h after freezing, the straws were removed from LN2, placed 30 sec in air and 30 sec into a 39°C water bath. The samples were added to 3 ml of sperm TL medium and centrifuged twice at 700 g for 6 min. After the second centrifugation the supernatant was removed and the sperm pellet was resuspended in 2 ml of sperm TL medium for analysis. The efficacies of two cryoprotective media (EYG and MG) on EP freezing were compared across three cool storage times. Two straws per testicle per cryoprotective media (total number of straws = 128) were used for analysis.

### Sperm analyses

An aliquot (5 µl) of the sperm suspension was added to 95 µl of sperm TL medium to make 1:20 sperm dilution. The percentage of motile sperm (on a 10 µl sample of the sperm dilution) was subjectively evaluated ( $\times 100$ ) in at least four separated fields on a warmed slide before freezing and immediately after thawing. At each time point an aliquot of the sperm dilution (20 µl) was mixed (1:1 v/v) with an aliquot (20 µl) of stain containing Eosin B/nigrosin, co-incubated for 1 min, smeared across a glass slide and quickly dried with a hair dryer. Stained slides were assessed (200 sperm/slide) using light microscopy (Nikon, Japan). Sperm with a pink or red like acrosome were considered nonviable. Sperm with clear or light acrosome color were considered viable.


### Statistical analysis

The percentage of sperm motility and viability were subjected to arcsin

transformation before analysis. The percentages of progressive motile and viable sperm within cool storage times were analyzed by General Linear Model of SAS (SAS, 1996). The effects of cool storage times, types of cryoprotective media and their interaction on sperm viability and progressive motility were examined using the General Linear Model of SAS with Tukey's multiple comparison test (SAS, 1996). Data was presented as least square means  $\pm$  SE of means.

### Results

Sperm viability (%) decreased (Fig. 1) after 24 and 48 h ( $76.5 \pm 5.3$  and  $67.9 \pm 5.8$ , respectively;  $P > 0.05$ ) of cool storage in comparison with the control ( $91.1 \pm 5.6$ ). The percentage of progressive motile sperm (0 h:  $73.4 \pm 3.4$ , 24 h:  $37.5 \pm 3.5$  and 48 h:  $27.1 \pm 3.2$ ;  $P < 0.05$ ) decreased across three times of cool storage (Fig. 1).



**Fig. 1: Effects of duration of buffalo (*Bubalus bubalis*) testis-epididymal cool storage (0 h: n = 10, 24 h: n = 10 and 48 h: n = 12) on sperm viability and progressive motility. (Different letters (AB) above bars for each variable (viability or progressive motility) indicate significant difference;  $P < 0.05$ )**

By extruding the effects of types of media and duration of cool storage, the results of the present study show an impact of cryopreservation on viability ( $> 50\%$  decline: Pre freeze:  $78.5 \pm 2.6$  versus post thaw:  $33.0 \pm 2.4$ ) and motility ( $> 80\%$  decline, Pre freeze:  $46.0 \pm 1.9$  versus Post thaw:  $7.9 \pm 2.6$ ) of river buffalo EP. However, MG was more efficient than EYG in protecting the post thaw sperm viability

( $P < 0.05$ ) across times of cool storage (Table 1).

Cryopreservation (MG or EYG; Table 2) severely affected post thaw sperm progressive motility (Table 2;  $P < 0.05$ ). Post thaw (for MG and EYG) progressive motility was significantly decreased across times of cool storage. Except for 48 h of cool storage, MG protected the sperm motility better ( $P < 0.05$ ) than EYG (Table 2).

**Table 1: Effects of duration of testis-epididymis cool storage<sup>†</sup> and cryoprotective media<sup>‡</sup> on viability (L.S. mean  $\pm$  SEM) of buffalo epididymal sperm**

Duration of storage (hours)	No. testis-epididymis	Post-thaw sperm viability (%)	
		MG	EYG
0	10	46.2 $\pm$ 3.4 <sup>Aa</sup>	39.63 $\pm$ 4.2 <sup>Ba</sup>
24	10	45.5 $\pm$ 4.3 <sup>Aa</sup>	15.8 $\pm$ 4.02 <sup>Bb</sup>
48	12	38.45 $\pm$ 3.3 <sup>Aa</sup>	12.5 $\pm$ 4.6 <sup>Bb</sup>

<sup>†</sup> 0, 24 and 48 hours. <sup>‡</sup>MG: Whole Milk-Glycerol; EYG: Whole Egg yolk-Tris-Citrate-Fructose-Glycerol. <sup>AB</sup>Values with different superscript within rows significantly differ ( $P < 0.05$ ). <sup>abc</sup>Values with different superscript within columns significantly differ ( $P < 0.05$ )

**Table 2: Effects of duration of testis-epididymis cool storage<sup>†</sup> and cryoprotective media<sup>‡</sup> on progressive motility (L.S. mean  $\pm$  SEM) of buffalo epididymal sperm**

Duration of storage (hours)	No. testis-epididymis	Post-thaw progressive motility (%)	
		MG	EYG
0	10	16.5 $\pm$ 2.6 <sup>Aa</sup>	11.6 $\pm$ 2.5 <sup>Ba</sup>
24	10	8.9 $\pm$ 2.6 <sup>Ab</sup>	5.4 $\pm$ 2.6 <sup>Bb</sup>
48	12	4.9 $\pm$ 2.3 <sup>Ab</sup>	2.4 $\pm$ 2.5 <sup>Ab</sup>

<sup>†</sup> 0, 24 and 48 hours. <sup>‡</sup>MG: Milk-Glycerol; EYG: Whole Egg yolk-Tris-Citrate-Fructose-Glycerol. <sup>AB</sup>Values with different superscript within rows significantly differ ( $P < 0.05$ ). <sup>abc</sup>Values with different superscript within columns significantly differ ( $P < 0.05$ )

## Discussion

Sperm progressive motility decreased significantly across times of storage. The results of this study demonstrated that viable sperm can be harvested from epididymes of river buffalo, at least after 48 h as well as 24 h of cool storage. Buffalo semen was diluted with different extenders and stored successfully at 4°C for at least 72 h (Sansone *et al.*, 2000). Motile EP has been collected after cool storage within epididymis or out of epididymis in different species (Braun *et al.*, 1994; Kikuchi *et al.*, 1998; Stilley *et al.*, 2000; Bruemmer *et al.*, 2002; Yu and Leibo, 2002; Nichi *et al.*, 2007). These findings are in agreement with the studies that reported motility and

viability of dog EP did not change up to Days 2 (Stilley *et al.*, 2000), 5 (Ponglowhapan *et al.*, 2006) or 8 (Bruemmer *et al.*, 2002) after cool storage within epididymis. Studies on Sika deer (*Cervus nippon*) (Hishinuma *et al.*, 2003) and Iberian red deer (*Cervus elaphus hispanicus*) (Soler *et al.*, 2005) EP showed that viable sperm can be isolated up to days 7 and 3 after cool storage, respectively, but sperm motility decreased the storage period continued in both of the studies. A study on cat EP preservation in tris or tris plus 10% egg yolk (Hermansson and Axner, 2007) at 4°C found that egg yolk-tris promoted the viability of cat EP compared to tris alone. In contrast to the present study, Bissett and Bernard, (2005) reported that only 30% of eland (*Taurotragus oryx*) EP survived at day 3 after epididymis cool storage, and Bartels *et al.* (1999) reported an increase in morphological changes in African buffalo (*Syncerus caffer*) EP with increasing duration of storage at 6°C.

Decreased sperm motility across storage times in the present study may relate to the postmortem changes in epididymal lumen. Prolonged postmortem preservation of epididymis resulted in a severe decrease in the pH of epididymal lumen in wild ruminant (Martinez-Pastor *et al.*, 2005). Pyknosis and release of epithelial cells in the lumen of cool stored epididymis were also reported (Hishinuma *et al.*, 2003). Therefore, the postmortem changes can occur even under cool storage and result in reduced pH of epididymal lumen. A strong positive correlation between pH of epididymal lumen and sperm motility was reported (Martinez-Pastor *et al.*, 2005).

Cryopreservation severely affected the viability and progressive motility of buffalo EP. These results are in agreement with the findings of a previous study on cryopreservation of African buffalo (*Syncerus caffer*) EP which reported a more than 50% decline in post thaw sperm viability (Lambrechts *et al.*, 1999). In contrast, some studies showed a good freezability of Sika deer (*Cervus nippon*) (Hishinuma *et al.*, 2003), dog (Ponglowhapan *et al.*, 2006), cat (Hermansson and Axner, 2007), ram (Kaabi *et al.*, 2003) and goat (Blash *et al.*, 2000)

EP. Tiplady *et al.* (2002) reported that cryopreservation of cooled equine EP induced a decrease in motility but not in viability from pre freeze to post thaw. On the other hand, comparing ejaculated semen, Raizada *et al.* (1990) reported a higher sensitivity of buffalo semen than cattle against cryopreservation damage. Galli *et al.* (1993) reported poor freezability of buffalo semen and explained it by lower phospholipid contents of buffalo sperm cell membrane.

The results of the present study indicated a superior effect of MG than EYG (20% egg yolk) on freezing of river buffalo EP. There is no report of using whole cow milk for cryopreservation of EP. However, lactose based diluent was more effective than citrate or tris based diluents to protect viability and motility of alpaca EP against freezing damage (Morton *et al.*, 2007). Adverse effects of 20% egg yolk in tris-citrate and Triladyl™ compared to 6% egg yolk have been reported for Spanish ibex (*Capra pyrenacia*) EP freezing (Santiago-Moreno *et al.*, 2006). The results of the present study are in contrast with the previous report of Herold *et al.* (2004) who compared Andromed® and Triladyl™ for freezing African buffalo (*Syncerus caffer*) EP. They concluded that freezability of African buffalo EP with Triladyl™ was better than Andromed® (Herold *et al.*, 2004), and suggested an appropriate equilibration time of equal to or more than 4 h for African buffalo (*Syncerus caffer*) EP freezing (Herold *et al.*, 2006). The very low sperm freezability (viability and progressive motility) with egg yolk based diluent in the present study can be related to the susceptibility of buffalo sperm to the egg yolk which was previously reported for buffalo semen cryopreservation (Galli *et al.*, 1993; Kumar *et al.*, 1993). Various levels of egg yolk in a tris based diluent for cryopreservation of buffalo semen showed that egg yolk beyond 5% did not improve post thaw sperm motility (Sahni and Mohan, 1990). Protective effects of egg yolk on preservation of the lipoprotein sheath of a sperm cell have been established (Kumar *et al.*, 1993). However, egg yolk stimulates the enzyme system of sperm to deaminate certain specific amino acids normally

present in egg yolk and yield hydrogen peroxide which is toxic to sperm during storage under aerobic conditions (Sahni and Mohan, 1990). These effects may enhance the toxicity of dead sperm, perhaps by providing substrate for lipid peroxidation (Shannon and Curson, 1972) and prolongation of storage by reducing pH (Santiago-Moreno *et al.*, 2006). It should be noted that as the yolk concentration increased in the diluent, the pH of the medium decreases (Shannon, 1972). Perhaps this is the reason for the depressing effects of high amounts (20%) of yolk on sperm motility reported in the present and other studies (Sansone *et al.*, 2000; Santiago-Moreno *et al.*, 2006). Shannon and Curson (1972) reported the depressing effects of enhanced levels of egg yolk (20%) on sperm viability during incubation in 5 and 25°C in combination with dead sperm on the sperm viability that is due to increased prooxide formation.

In summary, viable and motile buffalo EP can be isolated after cool storage within epididymis up to 48 h. Cryopreservation has an adverse effect on buffalo EP. Milk-7% glycerol diluent is superior to a 20% level of egg yolk plus glycerol on buffalo EP freezing output. More studies must be done to find the optimized protocol for cryopreservation of river buffalo EP.

## Acknowledgement

This project was financially supported by Shahid Chamran University of Ahvaz.

## References

- Barile, VL (2005). Improving reproductive efficiency in female buffaloes. *Livest. Prod. Sci.*, 92: 183-194.
- Bartels, P; Lubbe, K; Smith, RL and Godke, RA (1999). Morphological changes of caudal epididymal spermatozoa of African buffalo (*Syncerus caffer*) after storage at 6°C. *Theriogenology*. 51: 279.
- Bissett, C and Bernard, RTF (2005). The effect of prolonged cold storage of eland (*Taurotragus oryx*) cauda epididymides on the spermatozoa: possible implications for the conservation of biodiversity. *Theriogenology*. 63: 1592-1604.
- Blash, S; Melican, D and Gavin, W (2000).

- Cryopreservation of epididymal sperm obtained at necropsy from goats. *Theriogenology*. 54: 899-905.
- Braun, J; Sakai, M; Hochi, S and Oguri, N (1994). Preservation of ejaculated and epididymal stallion spermatozoa by cooling and freezing. *Theriogenology*. 41: 809-818.
- Bruemmer, JE; Reger, H; Zibinski, G and Squires, EL (2002). Effect of storage at 5°C on the motility and cryopreservation of stallion epididymal spermatozoa. *Theriogenology*. 58: 405-407.
- Fernández-Santos, MR; Estes, MC; Montoro, V; Soler, AJ and Garde, JJ (2006). Influence of various permeating cryoprotectants on freezability of Iberian red deer (*Cervus elaphus hispanicus*) epididymal spermatozoa: effects of concentration and temperature of addition. *J. Androl.*, 27: 734-745.
- Galli, A; Bornaghi, V; Balduzzi, D; Buttazzoni, L and Aleandri, R (1993). Sexual behaviour and semen quality relating to Italian buffalo. *Proceedings of 3rd World Buffalo Congress*. Vol. 1, Varna, Bulgaria. PP: 562-570.
- Hermansson, U and Axner, E (2007). Epididymal and ejaculated cat spermatozoa are resistant to cold shock but egg yolk promotes sperm longevity during cold storage at 4 degrees C. *Theriogenology*. 67: 1239-1248.
- Herold, FC; Aurich, JE and Gerber, D (2004). Epididymal sperm from the African buffalo (*Syncerus caffer*) can be frozen successfully with AndroMed<sup>(R)</sup> and with Triladyl but the addition of bovine seminal plasma is detrimental. *Theriogenology*. 61: 715-724.
- Herold, FC; de Haas, K; Colenbrander, B and Gerber, D (2006). Comparison of equilibration times when freezing epididymal sperm from African buffalo (*Syncerus caffer*) using Triladyl or AndroMed. *Theriogenology*. 66: 1123-1130.
- Hishinuma, M; Suzuki, K and Sekine, J (2003). Recovery and cryopreservation of sika deer (*Cervus nippon*) spermatozoa from epididymides stored at 4 degrees C. *Theriogenology*. 59: 813-820.
- Hori, T; Hagiuda, K; Endo, S; Hayama, A; Kawakami, E and Tsutsui, T (2005). Unilateral intrauterine insemination with cryopreserved caudal epididymal sperm recovered from refrigerated canine epididymides. *J. Vet. Med. Sci.*, 67: 1141-1147.
- Hori, T; Ichikawa, M; Kawakami, E and Tsutsui, T (2004). Artificial insemination of frozen epididymal sperm in beagle dogs. *J. Vet. Med. Sci.*, 66: 37-41.
- Kaabi, M; Paz, P; Alvarez, M; Anel, E; Boixo, JC; Rouissi, H; Herraiez, P and Anel, L (2003). Effect of epididymis handling conditions on the quality of ram spermatozoa recovered post-mortem. *Theriogenology*. 60: 1249-1259.
- Kikuchi, K; Nagai, J; Kashiwazaki, N; Ikeda, H; Noguchi, J; Shimada, A; Soloy, E and Kaneko, H (1998). Cryopreservation and ensuing *in vitro* fertilization ability of boar spermatozoa from epididymides stored at 4°C. *Theriogenology*. 50: 615-623.
- Kumar, S; Sahni, KL; Benjamin, BN and Mohan, G (1993). Effect of various levels of yolk on deep freezing and storage of buffalo semen in different diluters without adding glycerol. *Buff. J.*, 1: 79-85.
- Lambrechts, H; van Niekerk, FE; Coetzer, WA; Cloete, SWP and van der Horst, G (1999). The effect of cryopreservation on the survivability, viability and motility of epididymal African buffalo (*Syncerus caffer*) spermatozoa. *Theriogenology*. 52: 1241-1249.
- Martinez-Pastor, F; Guerra, C; Kaabi, M; Diaz, AR; Anel, E; Herraiez, P; de Paz, P and Anel, L (2005). Decay of sperm obtained from epididymes of wild ruminants depending on postmortem time. *Theriogenology*. 63: 24-40.
- Morton, KM; Bathgate, R; Evans, G and Maxwell, WMC (2007). Cryopreservation of epididymal alpaca (*Vicugna pacos*) sperm: a comparison of Citrate, Tris and Lactose based diluents, and pellets and straws. *Reprod. Fertil. Dev.*, 19: 792-796.
- Naderfard, HR and Qanemy, AW (1997). Buffalo breeding in Islamic Republic of Iran. *Proceedings of the 5th World Buffalo Congress*. Caserta, Italy. PP: 942-943.
- Nichi, M; Rijsselare, T; Van Soom, A; De Clercq, JBP; Goovaerts, IGF; Barnabe, VH and Bols, PEJ (2007). Effect of bull epididymis storage conditions on cryopreserved epididymal sperm *in vitro* fertility and lipid peroxidation status. 33rd Annual International Embryo Transfer Society. 7-9 Jan., Kyoto, Japan. *Reprod. Fertil. Dev.*, 19: 126.
- Parrish, JJ; Susho-Parrish, J; Winer, MA and First, NL (1988). Capacitation of bovine sperm by heparin. *Biol. Reprod.*, 38: 1171-1180.
- Ponglowhapan, S; Chatdarong, K; Sirivaidyapong, S and Lohachit, C (2006). Freezing of epididymal spermatozoa from dogs after cool storage for 2 or 4 days. *Theriogenology*. 66: 1633-1636.
- Raizada, BC; Sattar, A and Pandey, MD (1990). A comparative study of freezing buffalo semen in two diluters. In: Acharya, RM;

- Lokeshwar, RR and Kumar, AT (Eds.), *Proceedings of 2nd World Buffalo Congress*. New Delhi, India. International Buffalo Federation. Roma, Italy. PP: 66-74.
- Sahni, KL and Mohan, G (1990). Effect of various levels of yolk on viability of buffalo semen at 37, 5 and 196 degrees C. In: Acharya, RM; Lokeshwar, RR and Kumar, S (Eds.), *Proceedings of the 2nd World Buffalo Congress*. New Delhi, India. International Buffalo Federation. Roma, Italy. PP: 44-65.
- Salisbury, GW and Fuller, HK (1941). Preservation of bovine spermatozoa in yolk-citrate diluent and field results from its use. *J. Dairy Sci.*, 24: 905-910.
- Sansone, G; Nastri, MJF and Fabbrocini, A (2000). Storage of buffalo (*Bubalus bubalis*) semen. *Anim. Reprod. Sci.*, 62: 55-76.
- Santiago-Moreno, J; Toledano-Diaz, A; Pulido-Pastor, A; Dorado, J; Gómez-Brunet, A and López-Sebastián, A (2006). Effect of egg yolk concentration on cryopreserving Spanish ibex (*Capra pyrenaica*) epididymal spermatozoa. *Theriogenology*. 66: 1219-1226.
- SAS (1996). *SAS/STAT User's Guide*. Release 6.03 Edn., Statistical Analysis Systems Institute, Inc., Cary, NC., PP: 550-641.
- Shannon, P (1972). The effect of egg yolk level and dose rate of semen diluted in caprogen. *Proceedings of 7th International Congress on Animal Reproduction and Artificial Insemination*. Munich, Germany. Vol. 2, PP: 1440-1442. Cited by: Sansone, G; Nastri, MJF and Fabbrocini, A (2000). Storage of buffalo (*Bubalus bubalis*) semen. *Anim. Reprod. Sci.*, 62: 55-76.
- Shannon, P and Curson, B (1972). Toxic effect and action of dead sperm on diluted bovine semen. *J. Dairy Sci.*, 55: 614-620.
- Soler, AJ; Estes, MC; Fernández-Santos, MR and Garde, JJ (2005). Characteristics of Iberian red deer (*Cervus elaphus hispanicus*) spermatozoa cryopreserved after storage at 5 degrees C in the epididymis for several days. *Theriogenology*. 64: 1503-1517.
- Stilley, K; Pope, CE; Leibo, SP; Denniston, RS and Godke, RA (2000). Survival of canine epididymal sperm stored at 4°C in the testicles. *Theriogenology*. 53: 489 (abs.).
- Tiplady, CA; Morris, LHA and Allen, WR (2002). Stallion epididymal spermatozoa: pre-freeze and post-thaw motility and viability after three treatments. *Theriogenology*. 58: 225-228.
- Wani, NA; Nowshari, MA and Wernery, U (2005). Storage of camel (*Camelus dromedarius*) epididymal spermatozoa in different extenders and at different temperatures. *Biol. Reprod., Sp. Iss. SI.*, 234 (abs.).
- Yu, I and Leibo, SP (2002). Recovery of motile, membrane-intact spermatozoa from canine epididymides stored for 8 days at 4°C. *Theriogenology*. 57: 1179-1190.
- Zaugg, NL and Almquist, JO (1972). Motility of spermatozoa and control of bacteria in bovine semen diluents containing penicillin and neomycin or epicillin. *J. Dairy Sci.*, 56: 202-206.