



IJVR

ISSN: 1728-1997 (Print)
ISSN: 2252-0589 (Online)

Vol. 18

No. 4

Ser. No. 61

2017

**IRANIAN
JOURNAL
OF
VETERINARY
RESEARCH**



Review Article

Trend of toxocariasis in Iran: a review on human and animal dimensions

Zibaei, M.¹ and Sadjjadi, S. M.^{2,3*}

¹Department of Parasitology and Mycology, School of Medicine, Alborz University of Medical Sciences, Karaj, Iran; ²Department of Parasitology and Mycology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran; ³Basic Sciences in Infectious Diseases Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

*Correspondence: S. M. Sadjjadi, Department of Parasitology and Mycology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran, and Basic Sciences in Infectious Diseases Research Center, Shiraz University of Medical Sciences, Shiraz, Iran. E-mail: smsadjjadi@sums.ac.ir

(Received 7 Mar 2017; revised version 13 May 2017; accepted 30 May 2017)

Summary

One of the neglected soil and/or food-borne diseases with international public health importance is toxocariasis. Human cases are being increasingly reported from Asian, African, Oceania, European and the American countries. Hence, human toxocariasis (HT) is now considered as a major zoonosis with global and regional importance. In Iran, human and animal toxocariasis is an endemic disease with clinical and epidemiologic health problem aspects. Doubtless, understanding the epidemiology and the trend of this important parasitic disease and its affecting factors will provide the establishment of effective prevention and control programs. To better understand the trend of toxocariasis researches in Iran, this study was performed to analyze different aspects of this zoonotic disease including history, life cycle, species, human animals and environmental studies, diagnostic aspects and treatments to find out the gaps, including different aspects of clinical signs in human patients, new and specific recombinant antigens based on the native antigens, new diagnostic tools, especially rapid diagnostic tests, paratenic hosts status and new treatment procedures which is necessary to be investigated in the future studies on this important zoonotic disease.

Key words: Iran, Larva migrans, *Toxocara canis*, *Toxocara cati*, Toxocariasis

Introduction

Soil-transmitted nematodes including toxocariasis are neglected in the international public health importance in comparison with other helminthic diseases (Hotez and Wilkins, 2009). Toxocariasis is a serious zoonotic parasitic disease infecting invertebrate and vertebrate and human beings with high impact on public health worldwide. It is caused by a nematode belonging to genus *Toxocara*. A variety of animals such as dogs, cats, foxes, wolves, rodents, rats, pigeons, lambs, chickens and cattle show infection rates that may reach to 90% in some areas (Schantz, 1989). Human infection with toxocariasis was rare up until the last four decades when clinical cases were reported (Despommier, 2003). The present review article describes the lesser-known contributions of Iranian researchers to better understand the trend of toxocariasis researches analysis in Iran with emphasis on different aspects of this important zoonotic disease including history, life cycle, species, human, animals and environmental studies, diagnostic aspects and treatments to find out the gaps including investigations on the different aspects of clinical signs in human patients, making new and specific recombinant antigens based on the native antigens, new diagnostic tools especially rapid diagnostic tests, paratenic hosts status, new treatment procedures specifically non-

invasive procedures and finally registration of patients which is needed to be filled with new researches in the future on this important zoonotic and food-borne disease.

History

Animal toxocariasis has been reported during intestinal worms investigations of dogs in unpublished documents; however, the first published report about animal *Toxocara* is by Sadighian in 1969 who studied helminth parasites of stray dogs and jackals in Shahsavari area, Caspian region, Iran (Sadighian, 1969). In an extensive work on helminths in wild animals *Toxocara canis* and *Toxocara cati* were shown in the north of Iran (Dalimi and Mobedi, 1992). However, the chronological order of other studies has been reflected in Table 1.

Life cycle

A wide variety of animals, including mammalian can become infected with *Toxocara* species by ingesting eggs containing larvae from contaminated soil and the consumption of contaminated raw meat or liver (Despommier, 2003). In the paratenic or transport hosts, the larvae do not develop to maturity, but migrate for months throughout host organs such as liver, lungs, kidneys, heart, brain and retina before lodging within

Table 1: Prevalence of human toxocariasis based on reports in different regions of Iran

Province/regions, county	Author(s), year	Subjects	Clinical aspect ^β	Infected (%) / cases (No.)
Zabol	Farivar and Rafat, (1991)	Children under 5 years ^α	VLM	2
Chahar Mahal Bakhtiari	Yousefi <i>et al.</i> , (2000)	Children under 15 years	VLM	5.3
Shiraz	Sadjjadi <i>et al.</i> , (2000)	Children 6-13 years	VLM	25.6
Tehran	Rokni <i>et al.</i> , (2000)	Patients 3-52 years ^α	VLM	10
Hamedan	Fallah <i>et al.</i> , (2003)	Children under 10 years	VLM	5.3
Babol	Mekaniki, (2006)	Forty-year-old ^α	OLM	1
Kermanshah	Akhlaghi <i>et al.</i> , (2006)	Children 2-12 years	VLM	8.5
Shiraz	Zibaei <i>et al.</i> , (2008)	Child 6 years ^α	VLM	1
Zanjan	Nourian and Amiri, (2009)	Children referred to health centers and hospitals	VLM	2.7
Ahvaz	Alavi and Sephidgaran, (2009)	Children with chronic cough	VLM	34.5
Sari	Sharif <i>et al.</i> , (2010)	Schoolchildren	VLM	25.0
Ahvaz	Alavi <i>et al.</i> , (2011)	Schoolchildren 6-15 years	VLM	19.7
Khorramabad	Rafiee Alavi and Nayebzadeh, (2011)	Woman 57 years ^α	VLM	1
Khorramabad	Zibaei <i>et al.</i> , (2013)	Epileptic patients	NT	11.8
East-Azerbaijan	Garedaghi <i>et al.</i> , (2013)	Children	VLM	29.5
Khorramabad	Zibaei and Ghorbani, (2014)	Multiple sclerosis patients	NT	14.7
Shiraz	Zibaei <i>et al.</i> , (2014)	Child 6 years ^α	OLM	1
Zahedan	Ghorbani-Ranjbari <i>et al.</i> , (2015)	Students with cough	VLM	11.3
Isfahan	Hosseini-Safa <i>et al.</i> , (2015)	Children 5-15 years	VLM	1.4
Shiraz	Sarkari <i>et al.</i> , (2015)	Hyper eosinophilic individuals	VLM	2.0
Babol	Ebrahimifard <i>et al.</i> , (2015)	Adult eosinophilia individuals	VLM	23.5
Arak	Miladi <i>et al.</i> , (2016)	Individuals referred to lab diagnosis	VLM	4.2
Tabriz	Momeni <i>et al.</i> , (2016)	30 people	VLM	29.3
Tehran	Einipour <i>et al.</i> , (2016)	Uveitis patients	OLM	6.2
Mashhad	Berenji <i>et al.</i> , (2016)	Owners domestic cats and dogs	VLM	20.4
Arak	Mosayebi <i>et al.</i> , (2016)	Asthmatic children	VLM	1.8

^α Case report, and ^β VLM: Visceral larva migrans, OLM: Ocular larva migrans, and NT: Neurotoxocariasis

host tissues in state of arrested development. The larvae are not encysted but remain exposed to the host environment, absorbing nutrients across the nematode cuticle (Aziz *et al.*, 2007). The larvae can survive in tissue for several years despite vigorous host immunologic responses to parasite antigens (Maizels, 2013).

Toxocara species

Toxocara as a nematode is taxonomically included within the order of Ascaridida. Among a total of 21 species within the *Toxocara* genus, 2 are of significant public health concern, namely, *T. canis* and *T. cati*, for which dogs and cats are the definitive hosts, respectively. *T. canis* (Wener, 1782) and *T. cati* (Shrank, 1788) are the causative agents of toxocariasis in canidae, felidae and in humans worldwide. Molecular studies on cat nematodes in Shiraz showed that, the most prevalent one is *T. cati* (Mikaeili, 2013). *Toxocara vitulorum* (syn. *Neoascaris vitulorum*) (Goeze, 1782) is an ascarid that is frequently found in ruminants in sub-tropical regions. Its main hosts are cattle and buffalo in tropical and sub-tropical countries (Roberts, 1989; Tavassoli and Tadayon, 2000).

Human toxocariasis (HT)

Clinical cases

Human toxocariasis is a zoonotic disease caused by ingestion of embryonated eggs of *T. canis* and *T. cati*, which originate from the faeces of definitive hosts contaminating the environment and the consumption of contaminated raw and undercooked meat. Children in their first decade of life are prone to infection because of their geophagic behavior and mouthing of objects, which is linked to a higher risk of exposure at sandboxes or playgrounds contaminated with dogs and cats faeces.

Human infection is mostly asymptomatic but can be associated with severe clinical syndrome due to organ injury by migrating larvae (Glickman *et al.*, 1979). Depending on the organs affected and the specificity of the symptoms, the predominant clinical syndromes are classified as visceral larva migrans (VLM), ocular larva migrans (OLM), and common, neurologic, and covert toxocariasis (Magnaval *et al.*, 2001). Diagnosis of HT is traditionally based on a combination of clinical, serology and histopathological interpretation (Despommier, 2003). However, sensitivity is low, as biopsy material may not always contain the larvae. Serological examination, using *in vitro* obtained excretory-secretory (ES) proteins of the larvae, is the best laboratory-based option for diagnosis (Gold-Standard test), and is considered a useful predictor of *Toxocara* spp. infection when coupled to relevant clinical data (Smith *et al.*, 2009).

Visceral larva migrans

The most common, VLM, is a febrile disease of childhood, particularly affecting children between the age of one and five years old. Visceral larva migrans has been sporadically reported from different parts of Iran.

The first report on VLM in Iranian patients was reported by Jamalian (1976), followed by Farivar and Rafat (1991) who reported a 5-year-old boy admitted to a university hospital because of fever (two months), hepatomegaly, and weight loss. The serological examination was performed and the patient was strongly suspected to have suffered from visceral toxocariasis. A review of the clinical history of 10 VLM patients was published 8-years thereafter (Rokni *et al.*, 2000). Zibaei *et al.* (2008) presented a case of VLM in a 6-year-old child, which was diagnosed by sonographic and biopsy finding, and subsequently confirmed by enzyme-linked immunosorbent assay (ELISA) test.

Ocular larva migrans

Ocular abnormalities are a frequent complication of toxocariasis. Ocular toxocariasis is typically a monocular disease of young children. Patients with this disease present with chronic unilateral uveitis and a marked vitreous opacification that overlies a primary eosinophilic granuloma. The first report on OLM was published in 2006 and describes the clinical history of a 40-year-old man presenting with pain and visually impaired. The acute uveitis was accompanied by a granuloma in the eye-funds (Mekaniki, 2006). Following a gap of 8 years, one case of ocular toxocariasis in a 6-year-old boy suffering from clinical manifestation suggestive of toxocarial OLM was published (Zibaei *et al.*, 2014).

Exudative retinal detachment, posterior synechiae, and a cyclitic membrane may be present in the OLM. *Toxocara granuloma* is white, dome-shaped, and confined principally to the retina. Serological tests like ELISA with *Toxocara* infective larval ES antigen are the gold standard for diagnosis of ocular toxocariasis.

Covert toxocariasis

Serological studies in the past two decades indicated that HT is more prevalent in at least 3 main foci, in western and southern regions. These are, (i) the northwest areas (East Azerbaijan province); (ii) southwest (Khuzestan province), and (iii) south regions (Fars province) (Fig. 1).




Fig. 1: Approximate distribution of human toxocariasis and *Toxocara* species in Iran. The majority of currently known HT prevalence from provinces of Khuzestan (Ahvaz, #1), East Azerbaijan (Tabriz, #2), Fars (Shiraz, #3), Mazandaran (Sari, #4), Sistan and Baluchistan (Zahedan, #5), Kermanshah (Kermanshah, #6), and Hamadan (Hamadan, #7)

The first population study based on serological tests in Iran was reported by Sadjjadi *et al.* (2000) using sera of 436 clinically healthy children. In their sample, 25.6% of subjects tested positive for *Toxocara* excretory-secretory (TES) antigens. In another study, Yousefi *et al.*

(2000) demonstrated that 5.3% of children less than 15 years old possessed the IgG antibodies to TES. In a mild-scale seroepidemiological study, Garedaghi (2013) collected 336 sera from infants and tested for TES antibodies. Antibodies were confirmed in 99 individuals (29.5%). Several publications dealing with HT in Iran were identified (Akhlaghi *et al.*, 2006; Mekaniki, 2006; Fallah *et al.*, 2007; Alavi and Sefidgaran, 2009; Nourian and Amiri, 2009; Sharif *et al.*, 2010; Alavi *et al.*, 2011; Zibaei *et al.*, 2014; Ghorbani-Ranjbari *et al.*, 2015; Hosseini-Safa *et al.*, 2015; Momeni *et al.*, 2016) (Table 1).

Just as in other parasitic infections, direct demonstration is the only way to make definite diagnosis of visceral toxocariasis. However, it is difficult to find the larva in either tissue biopsies or autopsies due to its very small size. Based on improvements in the field of serology, diagnosis of VLM is usually made by detection of the specific antibodies to TES, along with clinical manifestations such as eosinophilia, eosinophilic pneumonia.

Neurological toxocariasis

The sites of central nervous system (CNS) invasion by *Toxocara* larvae include the spinal cord and brain. Neurotoxocariasis (NT) is a CNS manifestation of *Toxocara* infection that is influenced by multiple factors, such as the number of ingested eggs, previous exposure, and host genetic factors, all of which contribute to the complex pathogenesis of NT (Xinou *et al.*, 2003). Zibaei and Ghorbani (2014) conducted a case-control study and found a significant association between *Toxocara* seropositivity and multiple sclerosis. The results showed that from 68 multiple sclerosis patients who participated in the study, 14.7% had positive anti-*Toxocara* antibodies.

Epilepsy is considered an important health problem in the developing countries such as Iran. A possible association between toxocariasis and epilepsy has been hypothesized and *Toxocara* infection has been suggested as cofactor for epilepsy. In a study by Zibaei *et al.* (2013), frequency of *Toxocara* infection in epileptic patients was 11.8% (n=10 out of 85).

Animal toxocariasis

Toxocara infection in the dogs: *T. canis*

Published papers on animal toxocariasis in peer-reviewed journals have been reviewed (Table 2). The first paper of animal toxocariasis in Iran has been published by Sadighian (1969) in stray dogs (35%) and jackal (10%) of Caspian Sea region in Iran, although unpublished documents show earlier reports of animal toxocariasis in Iran (Table 1). Similarly, Mirzaian *et al.* (1972) reported that 16.6% of domestic dogs were infected with *T. canis* in Tehran.

In a study in Gilan and Mazandaran provinces on dogs, jackals, cats, badgers, foxes, and wild cat in the provinces, provided by the Provincial Institute of

Table 2: Prevalence of animals *Toxocara* spp. infection in different Iranian governorates according to reports

Governorates	Author(s), year	Animals	Species	Infected (%)
Tehran	Makarechian, (1955)	Dogs	<i>T. canis</i>	76.0
Tehran	Mobedi, (1968)	Dogs	<i>T. canis</i>	66.0
Mazandaran	Sadighian, (1969)	Stray dogs	<i>T. canis</i>	35.0
Mazandaran	Sadighian, (1969)	Jackals	<i>T. canis</i>	10.0
Tehran	Mirzayans, (1971)	Domestic cats	<i>T. cati</i>	31.4
Tehran	Mirzayans <i>et al.</i> , (1972)	Domestic dogs	<i>T. canis</i>	16.6
Gilan and Mazandaran	Arfaa, (1972)	Dogs	<i>T. canis</i>	35.0
Gilan and Mazandaran	Arfaa, (1972)	Jackal	<i>T. canis</i>	10.0
Mazandaran and Gilan	Dalimi and Mobedi, (1992)	Wild cats	<i>T. cati</i>	13.0
Mazandaran	Gholami <i>et al.</i> , (1999)	Dogs and Jackals	<i>T. canis</i>	13.3
Fars	Sadjjadi <i>et al.</i> , (2001)	Stray cats	<i>T. cati</i>	52.8
Fars	Mehrabani <i>et al.</i> , (2002)	Stray dogs	<i>T. canis</i>	3.9
Fars	Zibaei <i>et al.</i> , (2007)	Stray cats	<i>T. cati</i>	42.6
Mazandaran	Sharif <i>et al.</i> , (2007)	Stray cats	<i>T. cati</i>	60.0
West-Azerbaijan	Tavassoli <i>et al.</i> , (2008)	Dairy calves	<i>T. vitulorum</i>	8.1
Iran	Meshgi <i>et al.</i> , (2009)	Golden Jackal (<i>Canis aureus</i>)	<i>T. canis</i>	32.4
Isfahan	Arbabi and Hooshyar, (2009)	Stray cats	<i>T. cati</i>	13.3
Khorasan Razavi	Razmi, (2009)	Dogs	<i>T. canis</i>	17.9
North Iran	Daryani <i>et al.</i> , (2009)	Stray dogs	<i>T. canis</i>	60.0
Iran	Meshgi <i>et al.</i> , (2009)	Red fox (<i>Vulpes vulpes</i>)	<i>T. canis</i>	10.8
Zanjan	Esmailzadeh <i>et al.</i> , (2009)	Stray cats	<i>T. cati</i>	8.0
East-Azerbaijan	Shirzadi <i>et al.</i> , (2010b)	Persian cats	<i>T. cati</i>	20.0
Semnan	Eslami <i>et al.</i> , (2010)	Stray dogs	<i>T. canis</i>	22.0
East-Azerbaijan	Shirzadi <i>et al.</i> , (2010a)	Stray cats	<i>T. cati</i>	39.0
Ilam	Bahrani <i>et al.</i> , (2011)	Stray dogs	<i>T. canis</i>	36.6
Khorasan Razavi	Borji <i>et al.</i> , (2011)	Stray cats	<i>T. cati</i>	28.8
Tabriz	Garedaghi and Safar Mashaei, (2011)	Pet and stray dogs	<i>T. canis</i>	12.0
Golestan	Ghaemi <i>et al.</i> , (2011)	Persian leopard (<i>Panthera pardus saxicolor</i>)	<i>T. cati</i>	2.0 ^f
Ilam	Bahrani <i>et al.</i> , (2011)	Stray cats	<i>T. cati</i>	20.7
Isfahan	Pestehchian <i>et al.</i> , (2012)	Stray dogs	<i>T. canis</i>	6.3
Kerman	Mirzaei and Fooladi, (2012)	Owned dogs	<i>T. canis</i>	4.3
West-Azerbaijan	Tavassoli <i>et al.</i> , (2012)	Sheep and pet dogs	<i>T. canis</i>	9.7
Fars	Mikaeili <i>et al.</i> , (2013)	Stray cats	<i>T. cati</i>	26.7
Khorasan Razavi	Adinezadeh <i>et al.</i> , (2013)	Stray dogs	<i>T. canis</i>	7.3
Khorasan Razavi	Beironvand, <i>et al.</i> , (2013)	Domestic and stray dogs	<i>T. canis</i>	25.0
Khorasan Razavi	Shemshadi <i>et al.</i> , (2014)	Carnivores	<i>T. canis</i>	75.0
Tehran	Meshgi <i>et al.</i> , (2014)	Meshy cats	<i>T. cati</i>	52.7
East-Azerbaijan	Garedaghi <i>et al.</i> , (2014)	Stray dogs	<i>T. canis</i>	12.0
Hamadan	Sardarian <i>et al.</i> , (2015)	Household dogs	<i>T. canis</i>	41.8
Ardebil	Zare-Bidaki <i>et al.</i> , (2015)	Dogs	<i>T. canis</i>	43.5
Khorasan-Razavi	Emampour <i>et al.</i> , (2015)	Stray dogs	<i>T. canis</i>	29.0
Hamadan	Sardarian <i>et al.</i> , (2015)	Stray dogs	<i>T. canis</i>	22.1
Golestan	Geraili <i>et al.</i> , (2016)	Persian leopard	<i>T. canis</i>	23.3
East-Azerbaijan	Hajjipour <i>et al.</i> , (2016)	Stray cats	<i>T. cati</i>	78.8
Mazandaran	Savari <i>et al.</i> , (2016)	Dogs	<i>T. canis</i>	27.0
Golestan	Geraili <i>et al.</i> , (2016)	Persian leopard	<i>T. cati</i>	3.3

^f *T. cati* isolated from two leopards

Veterinary Medicine, a total sample size of 152 was assessed. Intestines were collected and direct examination was conducted. Parasite adults of *T. canis* were detected in dogs (6.2%) and jackals (15.5%) (Dalimi and Mobedi, 1992).

Another Iranian investigation on *T. canis* infection in dogs originates from 1992 and was a cross-sectional study conducted to determine the prevalence in stray dogs of Mazandaran province (Gholami *et al.*, 1999). Intestines of the animals were sampled from 75 randomly selected in each of the cities of Mazandaran province. Adult worms in the dogs' intestine were detected by stereomicroscope and direct microscope. The overall prevalence of infected dogs was 13.3% (n=10 out of 75).

A more recent study carried out in 2015 focused on the detection of intestinal helminths of stray dogs in Mashhad, in the northeast of Iran (Emampour *et al.*, 2015). In this study, the prevalence of *T. canis* was determined by the detection of adult worms in the intestine of animals, and not by the detection of eggs in

eggs in faeces. Adult worms of *T. canis* were detected in 24.0% of the dogs surveyed (n=29 out of 100). In accordance with the above survey, the significant differences in prevalence were not observed between male (n=14, 22.9%) and female (n=15, 38.4%). Moreover, the prevalence was higher in dogs younger's than 1 year (n=21, 25.6%) versus dogs below or above 1 year of age (n=8, 44.1%).

The reported data suggest that prevalence of infection in dogs, ranges from 4.3% to 43.5% in Iran, depending both on the method used and on the location. One of the best options for reducing toxocariasis in dogs is preventive chemotherapy (Dalimi and Mobedi, 1992). To be effective, this intervention should reach both stray and owner dogs.

Toxocara infection in the cats: *T. cati*

A large number of studies have been conducted for infection of definitive animal hosts with *T. cati*. In the case of feline infection, a variety of different methods have been used for identification of feline infection with

some measuring coproantigen positivity, some identifying worms following arecoline purgation, and some attempting to adjust these estimates in order to estimate the true infection status. However, the prevalence of infection with toxocariasis in cat in different regions of Iran is shown in Table 2 (Mirzayan, 1971; Arbabi and Hooshyar, 2009).

The early citation for feline toxocariasis in Iran was in 1971; it was mentioned that *T. cati* occurred in 31.4% of the domestic cats (Mirzayan, 1971). In 1992, a study in Gilan and Mazandaran provinces showed that the prevalence of toxocariasis in examined cats was 13.0% (Dalimi and Mobedi, 1992). The highest (78.0%) and lowest (8.0%) rate of infection were in cities of Azarshahr (East Azerbaijan province) and Zanjan (Zanjan province), respectively (Sharif *et al.*, 2007; Esmaeilzadeh *et al.*, 2009). Accordingly, all these animals having the label of final hosts can be regarded as the source of infection. The public health authorities have established a monitoring system to decrease the risk of infection.

Epidemiological picture of *Toxocara* spp. in paratenic host

Post-mortem examination (necropsy) of the tissues and organs for presence of larvae is the gold standard examination for the detection of infection in paratenic

hosts. Beaver (1969) hypothesized that organs or tissues of infected animals can serve as sources of *Toxocara* infection for humans.

In Iran, several studies have reported that gerbils, rats, chicken, and partridge as paratenic hosts infected with *Toxocara* spp. (Azizi *et al.*, 2007; Oryan *et al.*, 2009; Oryan *et al.*, 2010; Zibaei *et al.*, 2010d; Ebrahimi *et al.*, 2013; Zibaei *et al.*, 2017b) (Table 3). Azizi *et al.* (2007) experimentally inoculated *T. cati* eggs into the gizzard of chickens and observed migrating larvae in the liver. More recently, a study on naturally infected broiler chickens revealed that the frequencies of the species in the chickens were *T. canis* larvae (83.3%, n=5 out of 33) and *T. cati* larvae (16.7%, n=1 out of 33) (Zibaei *et al.*, 2017b).

Contamination with *Toxocara* species eggs in environment

Soil-transmitted and helminths are still one of the most important health problems in the world, even in developing countries (Alonso *et al.*, 2001). Although dogs and cats are definitive hosts, but expelled eggs should remain in the soil until larva develops within 6 weeks (Coelho *et al.*, 2001; Chorazy and Richardson, 2005). We found eleven reports conducted in Iran from 2006 (Tavassoli *et al.*, 2008; Maraghi *et al.*, 2014) (Table 4). The first Iranian study dealing with environmental

Table 3: *Toxocara* species identified in the paratenic hosts according to the technique applied

Host	Author(s), Year	Infection	Method	Species
Chickens	Oryan <i>et al.</i> , (2009)	Experimentally	D ^a	<i>T. cati</i>
Chickens	Oryan <i>et al.</i> , (2009)	Experimentally	D	<i>T. cati</i>
Chickens	Azizi <i>et al.</i> , (2007)	Experimentally	D	<i>T. cati</i>
Partridges	Ebrahimi <i>et al.</i> , (2013)	Naturally	D	-
Gerbils and rats	Zibaei <i>et al.</i> , (2010d)	Experimentally	D	<i>T. cati</i>
Chickens	Zibaei <i>et al.</i> , (2017b)	Naturally	D & M ^r	<i>T. canis</i> and <i>T. cati</i>

^a D: Digestion, and ^r M: Molecular

Table 4: Soil and vegetables contamination with eggs of *Toxocara* spp. in different regions of Iran

Area examined	Author(s), year	Subjects	Prevalence (%)
Shiraz	Motazedian <i>et al.</i> , (2006)	Public area	6.3
Urmia	Tavassoli <i>et al.</i> , (2008)	Public parks	7.8
Shiraz	Zibaei and Sadjjadi, (2010b)	Public area	Let. to Editor
Khorramabad	Zibaei <i>et al.</i> , (2010a)	Public parks	63.3
Tehran	Tavalla <i>et al.</i> , (2012)	Public parks	38.0
Tabriz	Garedaghi <i>et al.</i> , (2012)	Public parks	9.3
Tehran	Khazan <i>et al.</i> , (2012)	Public area	10.0
Qazvin	Saraei <i>et al.</i> , (2012)	Public parks	3.6
Southern Iran	Olyaei and Hajivandi, (2013)	Leafy vegetables plants	16.0
Amol	Siyadatpanah <i>et al.</i> , (2013)	Applied vegetables	3.2
Abadan	Maraghi <i>et al.</i> , (2014)	Public parks	29.2
Shiraz	Ghorbani Ranjbary <i>et al.</i> , (2014)	Public parks	15.0
Ahvaz	Khademvatan <i>et al.</i> , (2014)	Public area	46.3
Mashhad	Berenji <i>et al.</i> , (2015)	Public parks	7.7
Isfahan	Ghomashlooyan <i>et al.</i> , (2015)	Public parks	28.6
Khaf	Berenji <i>et al.</i> , (2015)	Public parks	10.3
Shahrekord	Fallah <i>et al.</i> , (2016)	Vegetables	3.9
Mazandaran	Rostami <i>et al.</i> , (2016)	Salad vegetables	1.7
Kermanshah	Ghashghaei <i>et al.</i> , (2016)	Public parks	18.0
Karaj	Zibaei <i>et al.</i> , (2017a)	Public parks	36.4

contamination detected *Toxocara* spp. eggs in 12 out of 112 soil samples collected by cluster random sampling from the uppermost centimeter of soil in 26 public places and children's playground in 4 regions of Shiraz, southern Iran, reported an overall prevalence of 6.3% with *T. cati* ova infected regions (Motazedian *et al.*, 2006). The highest contamination rate was in downtown public places in the 3rd (22.2%) and the 4th (20.0%) regions from four regions. Contamination was not observed during the dry seasons. However, due to their applied methodology, it was argued that the conclusions reported by them was not entirely supported by their results (Zibaei and Sadjjadi, 2010b).

Recently, the contamination level of public parks samples was assumed in the Mashhad and Khaf cities, in the northeast of Iran (Berenji *et al.*, 2015). Three-hundred and forty soil samples of Mashhad (n=195) samples and Khaf (n=145) parks samples, using simple random method were collected and examined with flotation technique and direct smear, and were evaluated using a light microscope. *Toxocara* spp. eggs were present in 7.7% and 10.3% of public parks of Mashhad and Khaf, respectively. The authorities of cities should pay considerable attention to disease control of these zoonotic infections. This can be achieved by increasing the hygiene education of people and pet owners and controlling the number of stray cats and dogs.

Diagnostic aspects

The diagnosis of *Toxocara* infection in human is mainly based on clinical, epidemiological, and laboratory information, which include imagining finding, blood tests, eosinophilia, immunoglobulin E level, and serological tests (Fillaux and Magnaval, 2013). Enzyme-linked immunosorbent assay and Western blotting are two types of tests that are available for the immunodiagnosis of toxocariasis, both using TES antigens (Magnaval *et al.*, 1991; Ishiyama *et al.*, 2009; Roldan and Espinoza, 2009; Maraghi *et al.*, 2012). It has long been considered that ES proteins secreted by nematodes hold the key to their success as parasites (Stirewalt, 1963). The first report by de Savigny (1975) demonstrated that *T. canis* larvae could be cultured for long periods of time, and that these culture supernatants contained antigens that were specific in diagnosing HT. The nature of TES was extensively studied by Maizels *et al.* (1984, 2000), and Maizels and Page (1990).

In Iran, research primarily at the Tehran University of Medical Sciences investigated the use of TES from second-stage larvae for detection of serum antibodies to *T. canis* (Jahani *et al.*, 1996). In a study by Maleky and Massoud (1997) ES antigen of *T. canis* and *T. cati* were demonstrated to have fractions with molecular weight between 30-120 kDa. Finding showed that the common fraction is 67 kDa. Zibaei *et al.* (2016) studied the ES antigens of *T. cati* larvae using ELISA and also Western blotting for serodiagnosis of HT and determined that the ELISA analyses using *T. cati* ES antigens has a good sensitivity (96.7%) compared to *T. canis* ES as antigens

for diagnosis HT. Electrophoretic analyses of *T. cati* ES antigens revealed a range of 20-150 kDa fractions and application of Western blotting based on 42 and 50 kDa fractions of ES antigens can be recommended for the acute diagnosis of toxocariasis. Production of monoclonal antibodies against *T. cati* has been reported by Zibaei *et al.* (2010c) for diagnosis of toxocariasis which verified that capture-ELISA is sensitive enough to detection of 5 ng/ml of *T. cati* antigen.

Although the antigenicity and specificity of TES is fairly high, cross-reaction to other parasite, especially nematode parasites have been observed (Yamasaki *et al.*, 2000). To overcome this problem, Zahabiun *et al.* (2015) cloned *T. cati* recombinant TES-120 (rTES-120), expressed and compared with its *T. canis* homolog in an IgG4-Western blotting. The findings showed that the diagnostic sensitivity and specificity of *T. cati*, rTES-120 were 70% and 100%, respectively, and about *T. canis* rTES-120 were 57.4% sensitivity and 94.4% specificity. The results revealed that the serodiagnosis sensitivity was 76% when using rTES-120 for both species considered.

Treatment

Anthelmintics eliminate worms in the intestines of definitive hosts. Piperazine, Pyrantel pamoate, Fenbendazole have been used for treatment of *T. canis* in dogs and Piperazine, Pyrantel pamoate, Selamectin, for *T. cati* infection in cats (Ballweber, 2001).

Although hypobiotic larvae are thought to be difficult to kill in dogs and cats, Selamectin has been used as a preventive against *Toxocara* in dogs and cats (Taylor *et al.*, 2016). Anyhow, the efficacy against larvae in different hosts, at different body sites, or at different stages of larval development, is still not completely understood (Oryan *et al.*, 2009). In paratenic hosts the larva tend to migrate to the brain as their final site which is difficult for drug exposure (Oryan *et al.*, 2008).

Patients with severe toxocariasis, particularly if there is CNS involvement, can be treated with systemic acting and larvicidal anthelmintics. Effective results have been reported with use of albendazole, mebendazole, fenbendazole, and diethylcarbamazine (Magnaval, 1995; Oryan *et al.*, 2009; Oryan and Alabadi, 2015). Most of these results are obtained from experimental studies on mice as one of the animal models where administration of higher doses of anthelmintic started directly after inoculation with *Toxocara* larvae and continued for several days (Overgaauw and van Knapen, 2013). The anthelmintic dose should be increased gradually over a period of days and covered by concomitant administration of steroids. The best results and a low rate of side effects in man are reported for mebendazole at a daily dose of 20-25 mg/kg for 3 weeks or albendazole 400 mg twice daily for 7 days administered concomitantly with prednisone 0.5-1.0 mg/kg daily in cases of visceral or ocular toxocariasis (Zibaei *et al.*, 2008; Zibaei *et al.*, 2014; Allahdin *et al.*, 2015).

The literature search on toxocariasis (caused by *T.*

canis and *T. cati*) in Iranian people yielded only a limited number of reports. Data from studies and case reports were predominantly conducted in Iran, hampering a reliable estimation of the importance of toxocariasis in Iranian people. Yet, the information suggests that *Toxocara* infection of the cats and dogs hosts, and environmental contamination with *Toxocara* species eggs is remarkable. Information from HT is less conclusive. However, data is limited and scattered across different fields and over time. There is a clear need for more new data including the investigations on the different clinical views in human patients, i.e. VLM, OLM, covert and neurological toxocariasis epidemiology, making new and specific recombinant antigens based on the native *Toxocara* species, new diagnostic tools, especially lateral flow tests or rapid diagnostic tests, paratenic hosts status especially those as food sources, new treatment procedures specifically non-invasive procedures and finally registration of patients in different health centers which need to be filled with new researches in the future on this important zoonotic and food-borne disease. Dedicated studies including human, animal and environmental health data should be conducted in different geographical areas. The outcome of such studies will allow policy makers to set priorities and design strategies, combining accurate surveillance and prevention of this zoonotic and food-borne disease.

Acknowledgement

This review was granted by the Office of Vice Chancellor for Research at Shiraz University of Medical Sciences, Grant No.: 95-01-106-14003.

References

- Adinezhadeh, A; Kia, EB; Mohebbali, M; Shojaee, S; Rokni, MB; Zarei, Z and Mowlavi, G (2013). Endoparasites of stray dogs in Mashhad, Khorasan Razavi Province, northeast Iran with special reference to zoonotic parasites. *Iran J. Parasitol.*, 8: 459-466.
- Akhlaghi, L; Ourmazdi, H; Sarafnia, A; Vaziri, S; Jadidian, K and Leghaii, Z (2006). An investigation on the toxocariasis seroprevalence in children (2-12 years-old) from Mahidasht area of Kermanshah Province (2003-2004). *Razi J. Med. Sci.*, 13: 41-48.
- Alavi, SM; Hosseini, SA; Rahdar, M; Salmanzadeh, SH and Nikkhuy, AR (2011). Determination of seroprevalence rate of *Toxocara canis* in 6-15 years aged rural and urban school children in Ahvaz. *Jundishapur Sci. Med. J.*, 10: 240-248.
- Alavi, SM and Sefidgaran, GH (2009). Prevalence of antibodies against *Toxocara canis* in children with chronic cough in urban and rural areas of Ahvaz city. *J. Med.*, 8: 181-188.
- Allahdin, S; Khademvatan, S; Raffei, A; Momen, A and Raffei, R (2015). Frequency of *Toxoplasma* and *Toxocara* sp. antibodies in epileptic patients, in south western Iran. *Iran J. Child Neurol.*, 9: 32-40.
- Alonso, JM; Stein, M; Chamorro, MC and Bojanich, MV (2001). Contamination of soil with eggs of *Toxocara* in a subtropical city in Argentina. *J. Helminthol.*, 75: 165-168.
- Arbabi, M and Hooshyar, H (2009). Gastrointestinal parasites of stray cats in Kashan, Iran. *Trop. Biomed.*, 26: 16-22.
- Arfaa, F (1972). Status of infections with larva migrans in Iran. *Iran J. Public Health*. 2: 168-174.
- Azizi, S; Oryan, A; Sadjjadi, SM and Zibaei, M (2007). Histopathologic changes and larval recovery of *Toxocara cati* in experimentally infected chickens. *Parasitol. Res.*, 102: 47-52.
- Bahrani, A; Doosti, A; Nahravanian, H; Noorian, AM and Asbchin, SA (2011). Epidemiological survey of gastrointestinal parasites in stray dogs and cats. *Aus. J. Basic Appl. Sci.*, 5: 1944-1948.
- Ballweber, LR (2001). *Veterinary parasitology: the practical veterinarian series*. 1st Edn., Oxford, UK, Butterworth-Heinemann. PP: 143-144.
- Beaver, PC (1969). The nature of visceral larva migrans. *J. Parasitol.*, 55: 3-12.
- Beiomvand, M; Akhlaghi, L; Fattahi Massom, SH; Meamar, AR; Motevalian, A; Oormazdi, H and Razmjou, E (2013). Prevalence of zoonotic intestinal parasites in domestic and stray dogs in a rural area of Iran. *Prev. Vet. Med.*, 109: 162-167.
- Berenji, F; Movahedi Rudy, AG; Rudy, AG; Fata, A; Tavassoli, M; Mousavi Bazaz, M and Salehi Sangani, G (2015). Soil contamination with *Toxocara* spp. eggs in public parks of Mashhad and Khaf, north east of Iran. *Iran J. Parasitol.*, 10: 286-289.
- Berenji, F; Pouryousef, A; Fata, A; Mahmoudi, M; Salehi, M and Khoshnegah, J (2016). Seroepidemiology study of toxocariasis in the owners of domestic cats and dogs in Mashhad, Northeastern Iran. *Iran J. Parasitol.*, 11: 265-268.
- Borji, H; Razmi, G; Ahmadi, A; Karami, H; Yaghfoori, S and Abedi, V (2011). A survey on endoparasites and ectoparasites of stray cats from Mashhad (Iran) and association with risk factors. *J. Parasit. Dis.*, 35: 202-206.
- Chorazy, ML and Richardson, DJ (2005). A survey of environmental contamination with ascarid ova, Wallingford, Connecticut. *Vector Borne Zoonotic Dis.*, 5: 33-39.
- Coelho, LM; Dini, CY; Milman, MH and Oliveira, SM (2001). *Toxocara* spp. eggs in public squares of Sorocaba, Sao Paulo state, Brazil. *Rev. Inst. Med. Trop. Sao Paulo.*, 43: 189-191.
- Dalimi, A and Mobedi, I (1992). Helminth parasites of carnivores in northern Iran. *Ann. Trop. Med. Parasitol.*, 86: 395-397.
- Daryani, A; Sharif, M; Amouei, A and Gholami, S (2009). Prevalence of *Toxocara canis* in stray dogs, Northern Iran. *Pak. J. Bio. Sci.*, 12: 1031-1035.
- de Savigny, DH (1975). *In vitro* maintenance of *Toxocara canis* larvae and a simple method for the production of *Toxocara* ES antigen for use in serodiagnostic tests for visceral larva migrans. *J. Parasitol.*, 61: 781-782.
- Despommier, D (2003). Toxocariasis: clinical aspects, epidemiology, medical ecology, and molecular Aspects. *Clin. Microbiol. Rev.*, 16: 265-272.
- Ebrahimi, M; Rouhani, S; Khazan, H and Mobedi, E (2013). Survey of *Toxocara* larva in hunted partridges' viscera. *Res. Med.*, 36: 200-204.
- Ebrahimi, SF; Fakhar, M; Sharif, M and Shirvani, SDN (2015). Seroprevalence of *Toxocara* infection among adult individuals with eosinophilia in Babol, 2013. *J. Mazandaran Univ. Med. Sci.*, 25: 355-362.
- Einipour, S; Jalusian, F; Ramezani, A; Abadi, A; Hajjiliani, F; Zeinali, S and Biderouni, FT (2016). Toxocariasis antibody titration in patients with uveitis. *Koomesh*. 17: 903-909.

- Emampour, RS; Borji, H and Naghibi, A** (2015). An epidemiological survey on intestinal helminths of stray dogs in Mashhad, North-East of Iran. *J. Parasit Dis.*, 39: 266-271.
- Eslami, A; Ranjbar-Bahadori, RH; Meshgi, B; Dehghan, M and Bokaie, S** (2010). Helminth infections of stray dogs from Garmsar, Semnan province, central Iran. *Iran J. Parasitol.*, 5: 37-41.
- Esmailzadeh, M; Shamsfard, M; Kazemi, A; Khalafi, SA and Altome, SA** (2009). Prevalence of protozoa and gastrointestinal helminthes in stray cats in Zanjan Province, north-west of Iran. *Iran J. Parasitol.*, 4: 71-75.
- Fallah, M; Azimi, A and Taherkhani, H** (2003). Seroprevalence of toxocariasis in children aged 1-9 years in Western Islamic Republic of Iran, 2003. *East. Mediterr. Health J.*, 13: 1073-1075.
- Fallah, AA; Makhtumi, Y and Pirali-Kheirabadi, K** (2016). Seasonal study of parasitic contamination in fresh salad vegetables marketed in Shahrekord, Iran. *Food Cont.*, 60: 538-542.
- Farivar, KH and Rafat, S** (1991). Visceral larva migrans (VLM): report of 2 cases. *Med. J. Islam. Repub. Iran.*, 2: 94-97.
- Fillaux, J and Magnaval, JF** (2013). Laboratory diagnosis of human toxocariasis. *Vet. Parasitol.*, 193: 327-336.
- Garedaghi, Y** (2013). Seroprevalence of toxocariasis in children in East-Azerbaijan Province, Iran. *Cukurova Med. J.*, 38: 581-586.
- Garedaghi, Y and Safar Mashaei, S** (2011). Prevalence of gastrointestinal helminthic infestation in pet and stray dogs in Tabriz (East-Azerbaijan Province). *Iran. J. Anim. Vet. Adv.*, 10: 1477-1479.
- Garedaghi, Y and Shabestari-Asl, SA** (2012). Contamination rate of *Toxocara* spp. eggs in the public parks of Tabriz city: a short report. *J. Rafsanjan Univ. Med. Sci.*, 11: 173-178.
- Garedaghi, Y; Shabestari-Asl, A and Ahmadi, S** (2014). Prevalence of gastrointestinal helminthes in stray dogs of Tabriz city, Iran. *Greener J. Bio. Sci.*, 4: 135-138.
- Geraili, A; Maroufi, Y; Dabirzadeh, M; Noormohammadi, H and Khoshshima Shahraki, M** (2016). A survey of gastrointestinal helminth of stray dogs in Zabol city, southeastern of Iran. *Arch. Razi Inst.*, 71: 57-60.
- Ghaemi, P; Sadr-Shirazi, N and Ghaemi, P** (2011). Study on Helminth parasites of Persian leopard (*Panthera pardus saxicolor* Pocock, 1927) in Golestan National Park and Biosphere Reserve of Iran. *World Appl. Sci. J.*, 13: 1471-1473.
- Ghashghaei, O; Khedri, J; Jahangiri-Nasr, F; Hadi Hashemi, S and Nourollahi Fard, SR** (2016). Contamination of soil samples of public parks with *Toxocara* spp. eggs in Kermanshah, Iran. *J. Fac. Vet. Med. Istanbul Univ.*, 42: 47-50.
- Gholami, S; Mobedi, E; Ziaee, H and Sharif, M** (1999). Intestinal helminth parasites in dog and jackal in inferent areas of Sari in the years 1371-72. *J. Mazandaran Univ. Med. Sci.*, 9: 1-12.
- Ghomashlooyan, M; Falahati, M; Mohaghegh, MA; Jafari, R; Mirzaei, F; Kalani, H; Sangani, GS and Azami, M** (2015). Soil contamination with *Toxocara* spp. eggs in the public parks of Isfahan city, central Iran. *Asian Pac. J. Trop. Dis.*, (Suppl. 1), 5: S93-95.
- Ghorbani-Ranjbar, A; Ghorbani-Ranjbar, N; Varzandian, S and Ghorbani-Ranjbar, Z** (2015). Frequency of anti-*Toxocara* antibodies in students with chronic cough and eosinophilia in a rural area of Zahedan, Iran. *Zahedan J. Res. Med. Sci.*, 17: 15-18.
- Ghorbani-Ranjbar, A; Shafie, A and Anamipor, AD** (2014). Frequency of *Toxocara* egg in public parks Shiraz/Iran. *J. Kerman Univ. Med. Sci.*, 21: 174-179.
- Glickman, LT; Schantz, PM and Cypess, RH** (1979). Epidemiological characteristics and clinical findings in patients with serologically proven toxocariasis. *Trans R. Soc. Trop. Med. Hyg.*, 73: 254-258.
- Hajipour, N; Imani Baran, A; Yakhchali, M; Banan Khojasteh, SM; Sheikhzade Hesari, F; Esmailnejad, B and Arjmand, J** (2016). A survey study on gastrointestinal parasites of stray cats in Azarshahr, (East Azerbaijan province, Iran). *J. Parasit. Dis.*, 40: 1255-1260.
- Hosseini-Safa, A; Mousavi, SM; Bagh-Badorani, MB; Ghatreh-Samani, MG; Mostafaei, S and Yousofi-Darani, H** (2015). Seroepidemiology of toxocariasis in children (5-15 yr Old) referred to the pediatric clinic of Imam Hossein Hospital, Isfahan, Iran. *Iran J. Parasitol.*, 10: 632-637.
- Hotez, PJ and Wilkins, PP** (2009). Toxocariasis: America's most common neglected infection of poverty and a helminthiasis of global importance? *PLoS Negl. Trop. Dis.*, 3: e400.
- Ishiyama, S; Ono, K; Rai, SK and Uga, S** (2009). Method for detecting circulating *Toxocara canis* antigen and its application in human serum samples. *Nepal Med. Coll. J.*, 11: 9-13.
- Jahani, MR; Massoud, J and Shojahsadati, SA** (1996). Preparation and purification of excretory-secretory antigens of *Toxocara canis* second stage larvae and its application in the diagnosis of infected people. *Kowsar Med. J.*, 1: 147-157.
- Jamalian, RV** (1976). Visceral larval migrans and a case report in Iran. *J. Iran Med. Council.* 10: 36-39.
- Khademvatan, S; Abdizadeh, R and Tavalla, M** (2014). Molecular characterization of *Toxocara* spp. from soil of public areas in Ahvaz southwestern Iran. *Acta Trop.*, 135: 50-54.
- Khazan, H; Khazaei, M; Seyedtabaei, SJ and Mehrabi, A** (2012). Prevalence of *Toxocara* spp. eggs in public parks in Tehran city, Iran. *Iran J. Parasitol.*, 7: 38-42.
- Magnaval, JF** (1995). Comparative efficacy of diethylcarbamazine and mebendazole for the treatment of human toxocariasis. *Parasitology.* 110: 529-533.
- Magnaval, JF; Faber, R; Maurieres, P; Charlet, JP and de Larrad, B** (1991). Application of the Western blotting procedure for the immunodiagnosis of human toxocariasis. *Parasitol. Res.*, 77: 697-702.
- Magnaval, JF; Glickman, LT; Dorchie, P and Morassin, B** (2001). Highlights of human toxocariasis. *Korean J. Parasitol.*, 39: 1-11.
- Maizels, RM** (2013). *Toxocara canis*: molecular basis of immune recognition and evasion. *Vet. Parasitol.*, 193: 365-374.
- Maizels, RM and Page, AP** (1990). Surface associated glycoproteins from *Toxocara canis* larval parasites. *Acta Trop.*, 47: 355-364.
- Maizels, RM; de Savigny, D and Ogilvie, BM** (1984). Characterization of surface and excretory-secretory antigens of *Toxocara canis* infective larvae. *Parasite Immunol.*, 6: 23-37.
- Maizels, RM; Tetteh, KK and Loukas, A** (2000). *Toxocara canis*: genes expressed by the arrested infective larval stage of a parasitic nematode. *Int. J. Parasitol.*, 30: 495-508.
- Makarechian, M** (1955). Study of intestinal helminths in dogs Tehran. DVM's Thesis. Tehran: Tehran University. PP: 15-80.
- Maleky, F and Massoud, J** (1997). Comparison of *Toxocara*

- canis* and *Toxocara cati* excretory-secretory proteins antigens second stage larvae by biochemical methods. *Pulse J.*, 6: 18-21.
- Maraghi, S; Mazhab Jafari, K; Sadjjadi, SM; Latifi, SM and Zibaei, M** (2014). Study on the contamination of Abadan public parks soil with *Toxocara* spp. eggs. *J. Environ. Health Sci. Eng.*, 12: 86.
- Maraghi, S; Rafiei, A; Hajhosseini, R and Sadjjadi, SM** (2012). Seroprevalence of toxocariasis in hyper eosinophilic individuals in Ahwaz, southwestern Iran. *J. Helminthol.*, 86: 241-244.
- Mehrabani, D; Sadjjadi, SM and Oryan, A** (2002). Prevalence of gastrointestinal nematode parasites in stray dogs in Shiraz, Southern Iran. *J. Appl. Anim. Res.*, 22: 157-160.
- Mekaniki, E** (2006). Ocular toxocariasis in a 40-year-old man. *J. Babol Univ. Med. Sci.*, 8: 67-69.
- Meshgi, B; Eslami, A; Bahonar, AR; Kharrazian-Moghadam, M and Gerami-Sadeghian, A** (2009). Prevalence of parasitic infections in the red fox (*Vulpes vulpes*) and golden Jackal (*Canis aureus*) in Iran. *Iran. J. Vet. Res.*, 10: 387-391.
- Meshgi, B; Jamshidi, S; Saadati, D; Hooshyar, H and Bokaie, S** (2014). An overview of *Toxocara cati* infection in stray cats in the metropolitan region of Tehran, Iran, and a comparison of two diagnostic methods. *Iran J. Vet. Med.*, 4: 53-56.
- Mikaeili, F; Mirhendi, H; Hosseini, M; Asgari, Q and Kia, EB** (2013). *Toxocara* nematodes in stray cats from Shiraz, southern Iran: intensity of infection and molecular identification of the isolates. *Iran J. Parasitol.*, 8: 593-600.
- Miladi, H; Didehdar, M; Hajhossein, R; Rafiei, F and Eslamirad, Z** (2016). Evaluating correlation between serum liver enzymes and toxocariasis: a case control study. *Asian Pac. J. Trop. Dis.*, 6: 460-462.
- Mirzaei, M and Fooladi, M** (2012). Prevalence of intestinal helminthes in owned dogs in Kerman city, Iran. *Asian Pac. J. Trop. Med.*, 5: 735-737.
- Mirzayans, A** (1971). Incidence of gastrointestinal helminths of domestic cats in the Tehran area of Iran. *J. Parasitol.*, 57: 1296.
- Mirzayans, A; Eslami, AH; Anwar, A and Sanjar, M** (1972). Gastrointestinal parasites of dogs in Iran. *Trop. Anim. Health Prod.*, 4: 58-60.
- Mobedi, I** (1968). Determine the extent of contamination of Tehran dogs to *Echinococcus granulosus*. MPH's Thesis, Tehran: Tehran University. . PP: 42-68.
- Momeni, T; Mahami Oskouei, M; Fallah, E; Safaiyan, A and Mahami Oskouei, L** (2016). Latent and asymptomatic *Toxocara* infection among young population in northwest Iran: the necessity of informing people as a potential health risk. *Scientifica (Cairo)*. 2016; 3562056. doi: 10.1155/2016/3562056.
- Mosayebi, M; Moini, L; Hajhossein, R; Didehdar, M and Eslamirad, Z** (2016). Detection of specific antibody reactivity to *Toxocara* larval excretory-secretory antigens in asthmatic patients (5-15 years). *Open Microbiol. J.*, 10: 162-167.
- Motazedian, H; Mehrabani, D; Tabatabaee, SH; Pakniat, A and Tavalali, M** (2006). Prevalence of helminth ova in soil samples from public places in Shiraz. *East. Mediterr. Health J.*, 12: 562-565.
- Nourian, A and Amiri, M** (2009). Seroepidemiology toxocariasis in children 2 to 15 years old referring to health centers and hospitals of Zanjan province in 1386. *J. Med.*, 47: 131-134.
- Olyaei, A and Hajivandi, L** (2013). Parasitological contamination of markets and farms in vegetables consumed in Southern Iran. *Global Vet.*, 10: 327-331.
- Oryan, A and Alidadi, S** (2015) Toxocariasis: a neglected parasitic disease with public health importance. *Trop. Med. Surg.*, 3: e126.
- Oryan, A; Sadjjadi, SM and Azizi, S** (2009). The effects of benzimidazoles on the larval stage of *Toxocara cati* in experimentally infected chickens. *Trop. Biomed.*, 26: 30-39.
- Oryan, A; Sadjjadi, SM and Azizi, S** (2010) Longevity of *Toxocara cati* larvae and pathology in tissues of experimentally infected chickens. *Korean J. Parasitol.*, 48: 79-80.
- Overgaauw, PA and van Knapen, F** (2013). Veterinary and public health aspects of *Toxocara* spp. *Vet. Parasitol.*, 193: 398-403.
- Pestehchian, N; Rasouli, A and Yousefi, HA** (2012). Distribution of intestinal worms among stray dogs in Isfahan, Iran. *J. Isfahan. Med. Sch.*, 29: 2827-2833.
- Rafiee Alavi, SE and Nayebzadeh, H** (2011). Presence of visceral larva migrans in the urinary bladder of a woman in Khorramabad, Iran. *Iran J. Parasitol.*, 6: 92-94.
- Razmi, GH** (2009). Survey of dogs' parasites in Khorasan Razavi province, Iran. *Iran J. Parasitol.*, 4: 48-54.
- Roberts, JA** (1989). The extraparastic life cycle of *Toxocara vitulorum* in the village environment of Sri Lanka. *Vet. Res. Commun.*, 13: 377-388.
- Rokni, MB; Massoud, J and Mowlavi, GH** (2000). Report of 10 cases of visceral larval migrans in Iran. *Iran J. Public Health*. 29: 61-66.
- Roldan, WH and Espinoza, YA** (2009). Evaluation of an enzyme-linked immunoelectrotransfer blot test for the confirmatory serodiagnosis of human toxocariasis. *Mem. Inst. Oswaldo Cruz.*, 104: 411-418.
- Rostami, A; Ebrahimi, M; Mehravar, S; Fallah Omrani, V; Fallahi, S and Behniafar, H** (2016). Contamination of commonly consumed raw vegetables with soil transmitted helminth eggs in Mazandaran province, northern Iran. *Int. J. Food Microbiol.*, 225: 54-58.
- Sadighian, A** (1969). Helminth parasites of stray dogs and jackals in Shahsavar area, Caspian region, Iran. *J. Parasitol.*, 55: 372-374.
- Sadjjadi, SM; Khosravi, M; Mehrabani, D and Oryan, A** (2000). Seroprevalence of *Toxocara* infection in school children in Shiraz, southern Iran. *J. Trop. Pediatr.*, 46: 327-330.
- Sadjjadi, SM; Oryan, A; Jalali, AR and Mehrabani, D** (2001). Prevalence and intensity of infestation with *Toxocara cati* in stray cats in Shiraz, Iran. *Veterinarski Arch.*, 71: 149-157.
- Saraei, M; Zakilo, M; Tavazoei, Y; Jahanihashemi, H and Shahnazi, M** (2012). Contamination of soil and grass to *Toxocara* spp. eggs in public parks of Qazvin, Iran. *Asian Pac. J. Trop. Biomed.*, 2: S1156-S1158.
- Sardarian, K; Maghsood, AH; Ghiasian, SA and Zahirnia, AH** (2015). Prevalence of zoonotic intestinal parasites in household and stray dogs in rural areas of Hamadan, Western Iran. *Trop. Biomed.*, 32: 240-246.
- Sarkari, B; Lari, M; Shafiei, R and Sadjjadi, SM** (2015). A comparative seroprevalence study of toxocariasis in hyper eosinophilic and apparently healthy individuals. *Arch. Pediatr. Infect. Dis.*, 3: e17911.
- Sarvi, S; Daryani, A; Sharif, M; Rahimi, MT; Azami, D; Marhaba, Z; Ahmadpour, E and Mizani, A** (2014). Domestic dog as a human health hazard in north of Iran. *J. Parasit. Dis.*, 40: 930-934.
- Schantz, PM** (1989). *Toxocara* larva migrans now. *Am. J.*

- Trop. Med. Hyg., 41: 21-34.
- Sharif, M; Daryani, A; Barzegar, G; Nasrollahi, M and Khalilian, A** (2010). Seroprevalence of toxocariasis in schoolchildren in northern Iran. Pak. J. Biol. Sci., 13: 180-183.
- Sharif, M; Nasrolahei, M; Ziapour, SP; Gholami, S; Ziaei, H; Daryani, A and Khalilian, A** (2007). *Toxocara cati* infections in stray cats in northern Iran. J. Helminthol., 81: 63-66.
- Shemshadi, B; Ranjbar-Bahadori, S and Jahani, S** (2014). Prevalence and intensity of intestinal helminths in carnivores and primates at Vakilabad Zoo in Mashhad, Iran. Comp. Clin. Pathol., 24: 384-391.
- Shirzadi, SH; Hashemzadeh-Farhang, H and Davoudi, J** (2010a). Study on infestation of stray cats in Tabriz area to *Toxocara cati*. Vet. Res., 6: 69-73.
- Shirzadi, SH; Hashemzadeh-Farhang, H; Shabestari-Asl, A and Garedaghi, Y** (2010b). Study on infestation of Persian cats in Tabriz area by *Toxocara cati*. Vet. Clin. Pathol., 3: 691-698.
- Siyadatpanah, A; Tabatabaei, F; Zeydi, AE; Spotin, A; Fallah-Omrani, V; Assadi, M; Moradi, S; Rostami, A; Memari, F and Hajjaliani, F** (2013). Parasitic contamination of raw vegetables in Amol, North of Iran. Arch. Clin. Infect. Dis., 8: e15983.
- Smith, H; Holland, C; Taylor, M; Magnaval, JF; Schantz, P and Maizels, R** (2009). How common is human toxocariasis? Towards standardizing our knowledge. Trends Parasitol., 5: 182-188.
- Stirewalt, MA** (1963). Chemical biology of secretions of larval helminths. Ann. N. Y. Acad. Sci., 113: 36-53.
- Tavalla, M; Oormazdi, H; Akhlaghi, L; Razmjou, E; Moradi Lakeh, M; Shojaee, S; Hadighi, R and Meamar, AR** (2012). Prevalence of parasites in soil samples in Tehran public places. Afr. J. Biotechnol., 11: 4575-4578.
- Tavassoli, M; Hadian, M; Charesaz, S and Javadi, S** (2008). *Toxocara* spp. eggs in public parks of Urmia city, West Azerbaijan Province Iran. Iran J. Parasitol., 3: 24-29.
- Tavassoli, M; Javadi, S; Firozi, R; Rezaei, F; Khezri, A and Hadian, M** (2012). Hair contamination of sheepdog and pet dogs with *Toxocara canis* eggs. Iran J. Parasitol., 7: 110-115.
- Tavassoli, M and Tadayon, K** (2000). Studies on *Toxocara vitulorum* infection of native and dairy calves in Urmia, Iran. J. Fac. Vet. Med. Univ. Tehran. 55: 71-73.
- Taylor, MA; Coop, RL and Wall, RL** (2016). *Veterinary Parasitology*. 4th Edn., Oxford, UK, John Wiley and Sons, Inc., P: 315.
- Xinou, E; Lefkopoulos, A; Gelagoti, M; Drevelegas, A; Diakou, A; Milonas, I and Dimitriadis, AS** (2003). CT and MR imaging findings in cerebral toxocaral disease. Am. J. Neuroradiol., 24: 714-718.
- Yamasaki, H; Araki, K; Lim, PK; Zasmy, N; Mak, JW; Taib, R and Aoki, T** (2000). Development of a highly specific recombinant *Toxocara canis* second-stage larva excretory-secretory antigen for immunodiagnosis of human toxocariasis. J. Clin. Microbiol., 38: 1409-1413.
- Yousefi, H; Avijgan, M; Taheri, AM; Zakeri, E and Massoud, J** (2000). A survey about toxocariasis in Chaharmahal va Bakhtiari province in first 6 month of 2000. Iran J. Infect. Dis. Trop. Med., 13: 16-20.
- Zahabiun, F; Sadjjadi, SM; Yunus, MH; Rahumatullah, A; Moghaddam, MH; Saidin, S and Noordin, R** (2015). Production of *Toxocara cati* TES-120 recombinant antigen and comparison with its *T. canis* homolog for serodiagnosis of toxocariasis. Am. J. Trop. Med. Hyg., 93: 319-325.
- Zare-Bidaki, M; Mobedi, I; Ahari, SS; Habibzadeh, S; Naddaf, S and Siavashi, M** (2010). Prevalence of zoonotic intestinal helminths of canids in Moghan Plain, north-western Iran. Iran J. Parasitol., 5: 42-51.
- Zibaei, M; Abdollahpour, F; Birjandi, M and Firoozeh, F** (2010a). Soil contamination with *Toxocara* spp. eggs in the public parks from three areas of Khorram Abad, Iran. Nepal Med. Coll. J., 12: 63-65.
- Zibaei, M; Bahadory, S; Cardillo, N and Khatami, AR** (2017a). Soil contamination with eggs of *Toxocara* species in public parks of Karaj, Iran. Int. J. Enteric Pathogs., 5: 45-48.
- Zibaei, M; Firoozeh, F; Bahrami, P and Sadjjadi, SM** (2013). Investigation of anti-*Toxocara* antibodies in epileptic patients and comparison of two methods: ELISA and Western Blotting. Epilepsy Res. Treat., 2013: 156815.
- Zibaei, M and Ghorbani, B** (2014). Toxocariasis and multiple sclerosis: a case-control study in Iran. Neurol. Asia. 19: 283-286.
- Zibaei, M and Sadjjadi, SM** (2010b). Prevalence of helminth ova in soil samples from public places in Shiraz. East Mediterr. Health J., 16: 580.
- Zibaei, M; Sadjjadi, SM; Geramizadeh, B and Firoozeh, F** (2008). Hepatic toxocariasis in a child: a case report from Shiraz, Southern Iran. Hepat Mon., 8: 310-312.
- Zibaei, M; Sadjjadi, SM; Ishiyama, S; Sarkari, B and Uga, S** (2010c). Production of monoclonal antibody against *Toxocara cati* second-stage larvae and its application for the detection of circulating antigens. Hybridoma (Larchmt). 29: 217-220.
- Zibaei, M; Sadjjadi, SM and Jahadi-Hosseini, SH** (2014). *Toxocara cati* larvae in the eye of a child: a case report. Asian Pac. J. Trop. Biomed., (Suppl. 1), 4: S53-55.
- Zibaei, M; Sadjjadi, SM and Maraghi, S** (2017b). The occurrence of *Toxocara* species in naturally infected broiler chickens revealed by molecular approaches. J. Helminthol., 91: 633-636.
- Zibaei, M; Sadjjadi, SM and Sarkari, B** (2007). Prevalence of *Toxocara cati* and other intestinal helminths in stray cats in Shiraz, Iran. Trop. Biomed., 24: 39-43.
- Zibaei, M; Sadjjadi, SM; Sarkari, B and Uga, S** (2016). Evaluation of *Toxocara cati* excretory-secretory larval antigens in serodiagnosis of human toxocariasis. J. Clin. Lab. Anal., 30: 248-253.
- Zibaei, M; Sadjjadi, SM and Uga, S** (2010d). Experimental *Toxocara cati* infection in gerbils and rats. Korean J. Parasitol., 48: 331-333.