

Effect of lead on thyroid function in sheep

Badiei, K.¹; Nikghadam, P.^{2*}; Mostaghni, K.¹
and Zarifi, M.³

¹Department of Clinical Sciences, School of Veterinary Medicine, Shiraz University, Shiraz, Iran; ²Graduated from School of Veterinary Medicine, Shiraz University, Shiraz, Iran and Department of Veterinary Medicine, School of Agriculture, Rasht Branch, Islamic Azad University, Rasht, Iran; ³Ph.D. Student in Large Animal Internal Medicine, Department of Clinical Sciences, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

*Correspondence: P. Nikghadam, Graduated from School of Veterinary Medicine, Shiraz University, Shiraz, Iran and Department of Veterinary Medicine, School of Agriculture, Rasht Branch, Islamic Azad University, Rasht, Iran. E-mail: pegah.nikghadam@gmail.com

(Received 27 Sept 2008; revised version 10 May 2009; accepted 18 May 2009)

Summary

The objective of this experiment was to determine the effect of long-term low-dose administration of lead (Pb) on thyroid function in sheep. In this experiment 10 healthy Iranian male sheep, aged about one-year-old were randomly allocated into two equal groups of control (n = 5) and experiment (n = 5). Both groups were kept under the same conditions of food and environment. The treatment group received lead acetate (5 mg/kg/day) orally for 8 weeks. Blood sampling of two groups was done on the same days, on days 0 (before receiving lead acetate in treatment group), 14, 28, 42, 56 and 70 after lead administration, every morning (8 to 9 a.m.). The function of the thyroid was evaluated by measuring the levels of serum thyroid hormones T3, T4, FT3, FT4 and TSH. The hepatic function in both groups was evaluated by measuring hepatic enzyme activities including alanin amino transferase (ALT), aspartate aminotransferase (AST), γ glutamyl transferase (GGT), and total bilirubin (TBIL). Serum levels of T3 (on days 14, 56 and 70) T4 (on days 14, 28, 42, 56 and 70), FT3 (on days 56 and 70), FT4 (on days 42, 56 and 70) and TSH (on days 14, 28, 42, 56 and 70) significantly decreased in lead-treated sheep when compared to the control group (P<0.05). Serum enzyme activities of ALT (on days 28 and 42), AST (on days 28, 42, 56 and 70) and TBIL (on days 28 and 70) in Pb treated sheep increased significantly in comparison with the control group (P<0.05). Our findings suggest that Pb exposed sheep can be at risk of low thyroid function.

Key words: Sheep, Lead, Thyroid function

Introduction

Lead (Pb) is considered to be one of the major environmental pollutants and has been incriminated as a cause of accidental poisoning in domestic animals more than any other substance (Casas and Sordo, 2006). Lead that contaminates the environment is largely air-borne but is redeposited by dust into soil and water and is taken up by or exists on the surface of plants which are grazed by livestock (Bolter *et al.*, 1975). Cattle, sheep, and horses are good indicators of pollution on vegetation (Debackere, 1983). One of the primary sources of lead contamination in air, soil and water is combustion of fuels containing lead additives. The main source of excess lead

intake for cattle was that in paint (Clapp, 2001) until the restrictions on the use of lead-base pigments in paints. Other sources of ingestible lead include storage battery plates, radiator repair, secondary smelters, putty, linoleum, asphalt roofing, engine oil, insecticide baits, and contaminated feeds (Buck, 1970; Christian and Tryphonas, 1971; Singh *et al.*, 1994; Yu, 2004; Casas and Sordo, 2006). Lead can cause profound hematological, neurological, gastrointestinal, renal, rheumatological and endocrine manifestations in man even at levels previously considered safe (Cullen *et al.*, 1983; Pagliuca *et al.*, 1990; Lyons and Pahwa, 2005). Lead exposure also causes functional impairment of pituitary-adrenal axis as well as the pituitary-thyroid axis

(Singh and Dhawan, 1999). There are conflicting reports in the literature regarding the effect of occupational lead exposure on the thyroid functions in experimental animals and humans (Singh *et al.*, 2000). The thyroid gland plays an important role in energy usage, synthesis of RNA protein, consumption of oxygen by cells, overall body metabolism, growth processes and neurological development (Vanderpump and Tunbridge, 2008). In adult sheep, poor wool growth, depressed milk yield, reduced weight gain, impaired reproductive performance with loss of libido in rams and late abortions or birth of weak lambs with visibly enlarged thyroid glands as well as an increased susceptibility to infectious agents are the most prominent clinical findings in hypothyroid cases (Sipos *et al.*, 2004). As there was a knowledge gap on the effect of Pb on the thyroid function in sheep, the present study was designed to provide some data on this issue.

Materials and Methods

This study was performed under regulation of Shiraz University for using animals in scientific procedures. In this experiment 10 healthy male sheep, aged about one-year-old were randomly allocated into two equal groups of control ($n = 5$) and experiment ($n = 5$). The two groups were kept under the same conditions of food and environment. Water and hay were supplied *ad libitum* to all animals. Before commencing the experiment, samples of blood and faeces were submitted for hematological, bio-chemical and parasitological examinations to monitor the health status of the animals. The treatment group received 5 mg/kg/day of lead acetate orally for 8 consecutive weeks. Blood sampling of both groups was done on days 0 (before receiving lead acetate in treatment group), 14, 28, 42, 56 and 70 after lead administration, every morning (8 to 9 a.m.). Blood samples were collected by jugular veinipuncture in sterile silicone-coated vacutainers, allowed to clot, and then centrifuged at 3000 g for 10 min. Serum was separated and stored at -20°C until use. Serum T3, T4, FT3, FT4 and TSH were

measured by RIA (Bruss, 1997). Serum samples were digested and Pb was measured by atomic absorption spectrophotometry (Shimadzu AA-670; Shimadzu Corp, Kyoto, Japan) at a wavelength of 283.3 nm (Szkoda and Żmudzki, 2005). All glassware was acid washed with 2 M nitric acid. Standards were prepared from a certified Pb standard solution. Alanine aminotransferase (ALT) and aspartate aminotransferase (AST) activities were measured by modified Reitman-Frankel method. Serum GGT concentration was measured by Kinetic colorimetric test according to Szasz/Persijn (Szasz *et al.*, 1974). Total bilirubin was measured using Zist Chimi Kits (Zist Chimi, Iran). Data were checked for normality before performing analysis. All analyses utilized parametric statistical methods. Results were expressed as mean \pm SE. All data were analysed using SPSS for Windows, Version 15.0. In order to compare the mean control values of different parameters with the mean values of the same parameters measured in the treatment group on days 0, 14, 28, 42, 56 and 70 of Pb administration, Repeated measure ANOVA and Bonferroni post hoc test were used. The level of significance for each test was at $P < 0.05$.

Results

The results are indicated in Table 1 and 2. Serum Pb concentration (on different sampling days following lead administration) and serum enzyme activities of alanin amino transferase (ALT) (on days 28 and 42), aspartate aminotransferase (AST) (on days 28, 42, 56 and 70) and total bilirubin (TBIL) concentration (on days 28 and 70) increased significantly in lead treated sheep when compared with the control values ($P < 0.05$) (Table 1). No significant changes were observed in γ glutamyl transferase (GGT) activity in the treatment group when compared with the control values. Serum T3 (on days 14, 56 and 70), FT3 (on days 56 and 70) (Fig. 1), T4 (on days 14, 28, 42, 56 and 70), FT4 (on days 42, 56 and 70) (Fig. 2) and TSH (on days 14, 28, 42, 56 and 70) (Fig. 3) concentrations decreased significantly in the

Table 1: Concentrations (mean ± SE) of serum Pb and total bilirubin and serum hepatic enzyme activities before and after lead acetate administration on different days in sheep (n = 10)

Sampling days	0	14	28	42	56	70
Pb treatment (μmol/L)	0.00 ± 0.00	1.60 ± 0.60*	1.27 ± 0.36*	1.92 ± 0.06*	1.23 ± 0.21*	0.66 ± 0.25*
Pb control (μmol/L)	0.13 ± 0.13	0.18 ± 0.09	0.00 ± 0.00	0.03 ± 0.02	0.00 ± 0.00	0.00 ± 0.00
ALT treatment (IU/L)	39 ± 1.70	40 ± 3.83	61 ± 5.17*	141 ± 7.47*	33 ± 5.00	30 ± 4.37
ALT control (IU/L)	38 ± 2.34	41 ± 3.24	39 ± 1.93	34 ± 2.40	39 ± 4.50	38 ± 2.35
AST treatment (IU/L)	104 ± 1.70	110 ± 14.42	83 ± 6.60*	141 ± 7.50*	127 ± 9.12*	84 ± 7.33*
AST control (IU/L)	103 ± 3.05	95 ± 4.60	109 ± 8.51	117 ± 6.35	93 ± 3.57	103 ± 3.43
GGT treatment (IU/L)	60 ± 2.60	58 ± 7.02	70 ± 4.90	63 ± 7.50	48 ± 3.67	84 ± 4.53
GGT control (IU/L)	61 ± 3.32	52 ± 3.51	78 ± 4.62	59 ± 5.46	57 ± 5.50	61 ± 3.40
TBIL treatment (mg/dl)	0.37 ± 0.02	0.47 ± 0.07	0.64 ± 0.05*	0.37 ± 0.03	0.24 ± 0.05	0.19 ± 0.03*
TBIL control (mg/dl)	0.36 ± 0.02	0.36 ± 0.04	0.34 ± 0.03	0.40 ± 0.02	0.36 ± 0.03	0.37 ± 0.02

* Shows significant difference between treatment and control groups (P<0.05)

Table 2: Concentrations (mean ± SE) of thyroid hormones before and after lead acetate administration on different days in sheep (n = 10)

Sampling days	0	14	28	42	56	70
T3 treatment (nmol/L)	1.12 ± 0.11	0.84 ± 0.10*	0.92 ± 0.06	0.83 ± 0.04	0.60 ± 0.06*	0.74 ± 0.04*
T3 control (nmol/L)	1.13 ± 0.11	1.33 ± 0.17	1.06 ± 0.11	1.06 ± 0.18	1.08 ± 0.08	1.13 ± 0.11
T4 treatment (nmol/L)	83 ± 4.90	35 ± 3.00*	42 ± 4.55*	40 ± 2.92*	40 ± 5.00*	50 ± 6.14*
T4 control (nmol/L)	83 ± 4.81	88 ± 5.05	87 ± 7.03	83 ± 5.00	76 ± 5.05	83 ± 4.61
FT3 treatment (pmol/L)	5.94 ± 0.50	3.22 ± 0.37	4.06 ± 0.78	3.60 ± 0.40	3.20 ± 0.57*	3.15 ± 0.51*
FT3 control (pmol/L)	5.98 ± 0.43	6.75 ± 0.51	5.90 ± 0.45	6.10 ± 0.35	5.08 ± 0.60	5.97 ± 0.43
FT4 treatment (pmol/L)	30 ± 1.15	23 ± 2.24	28 ± 1.70	18 ± 1.80*	19 ± 1.70*	20 ± 2.33*
FT4 control (pmol/L)	30 ± 1.03	29 ± 1.45	32 ± 2.70	31 ± 2.45	27 ± 2.58	30 ± 1.02
TSH treatment (mIU/L)	0.13 ± 0.01	0.03 ± 0.00*	0.02 ± 0.00*	0.02 ± 0.00*	0.02 ± 0.00*	0.03 ± 0.00*
TSH control (mIU/L)	0.13 ± 0.01	0.11 ± 0.02	0.14 ± 0.02	0.14 ± 0.01	0.14 ± 0.02	0.13 ± 0.01

* Shows significant difference between treatment and control groups (P<0.05)




Fig. 1: Serum T3 and FT3 concentrations of control and Pb treated sheep




Fig. 2: Serum T4 and FT4 concentrations of control and Pb treated sheep




Fig. 3: Serum TSH concentration of control and Pb treated sheep

Pb treated group when compared with the control values (P<0.05) (Table 2).

Discussion

Serum levels of thyroid hormones, including T3, T4 and TSH, are commonly used as reliable indicators of the thyroid function in humans and experimental animals (Paier *et al.*, 1993; Chaurasia *et al.*, 1996; Kelly, 2000). All reactions necessary for the formation of T3 and T4 are influenced and controlled by pituitary

thyroid-stimulating hormone (TSH) which stimulates follicular cells in the thyroid gland (Beers and Berkow, 1999; Kelly, 2000). Pituitary TSH secretion is controlled by a negative feedback mechanism modulated by the circulating level of free T₄ and free T₃ and by conversion of T₄ to T₃ in the pituitary thyrotropic cells. T₃ is the metabolically active iodothyronine. TSH secretion is also influenced by thyrotropin-releasing hormone (TRH), a 3-amino acid peptide synthesized in the hypothalamus which stimulates the pituitary to release TSH (Beers and Berkow, 1999; Kelly, 2000; Higgins, 2007). In our study mean serum concentrations of T₃ (on days 14, 56 and 70), T₄ (on days 14, 28, 42, 56 and 70), FT₃ (on days 56 and 70), FT₄ (on days 42, 56 and 70) and TSH (on days 14, 28, 42, 56 and 70) decreased significantly in lead-treated sheep. Hypothyroidism occurring in human subjects with occupational exposure to lead is suggestive of a negative effect of the element on thyroid function (Lasisz *et al.*, 1992). Also Singh and Dhawan (1999) showed that lead treatments on rats at a dose rate of 50 mg/kg body weight for a period ranging from 1-4 months caused a significant depression in T₃ and T₄ levels. On the other hand, a dose related depression of thyroid functions had been observed in humans during occupational exposure to inorganic lead (Gustafson *et al.*, 1989). It has been proposed that Pb can accumulate in the organism and its actions include damage to cell membranes and disorders of the oxido-reductive processes in the cells (Lasisz *et al.*, 1992). Therefore thyroid dysfunction in our study might be related to structural damage of thyroid follicular cells due to accumulation of Pb in the thyroid gland. As the thyroid gland is the only organ involved in T₄ synthesis (Kelly, 2000), the decrease in the serum level of this hormone in the Pb treated sheep may suggest that Pb-induced thyroid dysfunction is mediated by influencing the production and/or secretion of T₄ by the follicular cell of the thyroid gland. Although on different days following lead administration free thyroid hormones reduced in our Pb exposed sheep, TSH didn't respond to low thyroid hormones and conversely significantly decreased. The significant decrease of serum TSH on

different days following lead administration in Pb treated sheep may also be the result of Pb effect on regulatory enzymes associated with hypothalamic pituitary thyroid (HPT) axis. On the other hand, thyroid hormones are metabolized in peripheral tissue by deiodination, conjugation, deamination, and decarboxylation enzyme reactions. Therefore, alterations in these metabolic pathways may significantly affect thyroid hormone metabolites and influence thyroid function at the cellular level (Paier *et al.*, 1993; Chaurasia *et al.*, 1996; Kelly, 2000). Most of the circulating T₃ originates from the extra-thyroidal tissues. The peripheral deiodination of T₄ to T₃, taking place mainly in the liver, is dependent on 5'-monodeiodinase (5'-D) activity (Paier *et al.*, 1993; Chaurasia *et al.*, 1996; Gibney *et al.*, 2003). A common feature of thyroid hormone dysfunction at cellular levels is a low level of circulating T₃, with generally normal to slightly elevated blood T₄ levels and either normal or slightly suppressed TSH levels (Kelly, 2000). Significant increase in serum enzyme activities of ALT, AST and TBIL in Pb treated sheep in our study could be an indication of hepatic cells damage (Latimer *et al.*, 2003). Hepatic pathology influence serum thyroid hormone concentrations, secondary to the effects on peripheral enzyme pathways (Kelly, 2000). Therefore, partial decrease of serum T₃ concentration in Pb treated sheep might be related to hepatic dysfunction. In conclusion, our study shows that long-term low-dose exposure to lead in sheep affects thyroid function due to hypothyroidism.

Acknowledgement

This work was sponsored by a grant from Shiraz University, Iran, No. 84-VE-1784-C308.

References

- Beers, MH and Berkow, R (1999). *The Merck manual of diagnosis and therapy*. 17th Edn., USA, John Wiley and Sons. P: 82.
- Bolter, E; Butz, TR and Arseneau, JF (1975). *Heavy metal mobilization by natural organic acids*. International conference on heavy metals in the environment. Toronto, Canada.

- P: C-81.
- Bruss, ML (1997). *Clinical biochemistry of domestic animals*. 1st Edn., New York, Academic Press. PP: 571-588.
- Buck, WB (1970). Lead and organic pesticide poisonings in cattle. *J. Am. Vet. Med. Assoc.*, 156: 1468-1472.
- Casas, JS and Sordo, J (2006). *Lead: chemistry, analytical aspects, environmental impact and health effects*. 1st Edn., Amsterdam, Elsevier Science, P: 19.
- Chaurasia, SS; Gupta, P; Kar, A and Maiti, PK (1996). Free radical mediated membrane perturbation and inhibition of type-I iodothyronine 5'-monodeiodinase activity by lead and cadmium in rat liver homogenate. *Biochem. Mol. Biol. Int.*, 39: 765-770.
- Christian, RG and Tryphonas, L (1971). Lead poisoning in cattle: brain lesions and hematologic changes. *Am. J. Vet. Res.*, 32: 203-216.
- Clapp, J (2001). *Toxic exports: the transfer of hazardous wastes from rich to poor countries*. 1st Edn., London, Cornell University Press. P: 65.
- Cullen, MR; Robins, JM and Eskenazi, B (1983). Adult inorganic lead intoxication: presentation of 31 new cases and a review of recent advances in the literature. *Medicine*. 62: 221-247.
- Debackere, M (1983). Environmental pollution: the animal as source, indicator, and transmitter. In: *Veterinary pharmacology and toxicology*. 2nd Edn., Westport, CT, AVI Publishing Co., PP: 595-608.
- Gibney, MJ; MacDonald, IA and Roche, HM (2003). *Nutrition and metabolism (The Nutrition Society Textbook)*. 1st Edn., Oxford, Blackwell Science. P: 81.
- Gustafson, A; Hedner, P; Schutz, A and Skerfving, S (1989). Occupational lead exposure and pituitary function. *Int. Arch. Occup. Environ. Health*. 61: 277-281.
- Higgins, C (2007). *Understanding laboratory investigations for nurses and health professionals*. 2nd Edn., Oxford, Blackwell Science. P: 142.
- Kelly, GS (2000). Peripheral metabolism of thyroid hormones: a review. *Altern. Med. Rev.*, 5: 306-333.
- Lasisz, B; Zdrojewicz, Z and Marcinkowski, Z (1992). Effect of lead on thyroid function. *Wiad. Lek.*, 45: 116-119.
- Latimer, KS; Mahaffey, EA and Prasse, KW (2003). *Duncan and Prasse's veterinary laboratory medicine: clinical pathology*. 4th Edn., Oxford, Blackwell Science. P: 194.
- Lyons, KE and Pahwa, R (2005). *Handbook of essential tremor and other tremor disorders*. 1st Edn., Boca Raton, FL, Taylor and Francis Group. P: 22.
- Pagliuca, A; Mufti, GJ; Baldwin, D; Lestas, AN; Wallis, RM and Bellingham, AJ (1990). Lead poisoning: clinical, biochemical, and haematological aspects of a recent outbreak. *J. Clin. Pathol.*, 43: 277-281.
- Paier, B; Hagemüller, K; Noli, MI; Gonzalez Pondal, M; Stiegler, C and Zaninovich, AA (1993). Changes induced by cadmium administration on thyroxine deiodination and sulfhydryl groups in rat liver. *J. Endocrinol.*, 138: 219-224.
- Singh, B; Chandran, V; Bandhu, HK; Mittal, BR; Bhattacharya, A; Jindal, SK and Varma, S (2000). Impact of lead exposure on pituitary-thyroid axis in humans. *Biometals*. 13: 187-192.
- Singh, B and Dhawan, D (1999). Effect of lithium on thyroidal ¹³¹iodine uptake, its clearance, and circulating levels of triiodothyronine and thyroxine in lead-treated rats. *Radiat. Environ. Biophys.*, 38: 261-266.
- Singh, B; Dhawan, D; Chand, B and Mangal, PC (1994). Biokinetics of iodine-131 in rat thyroid following lead and lithium supplementation. *Biol. Trace Elem. Res.*, 40: 287-293.
- Sipos, W; Miller, I; Fountoulakis, M; Schmall, F; Patz, M; Schwendenwein, I; Rapp, E; Taxacher, A and Gemeiner, M (2004). Hypothyroid goitre in a ram: chemical analysis gives indirect evidence for a structurally altered type of ovine thyroglobulin. *J. Vet. Med.*, 51(A): 90-96.
- Szasz, G; Weimann, G; Stahler, F; Wahlefeld, AW and Persijn, JP (1974). New substrates for measuring gamma-glutamyl transpeptidase activity. *Z. Klin. Chem. Klin. Biochem.*, 12: 228.
- Szkoda, J and Żmudzki, J (2005). Determination of lead and cadmium in biological material by graphite furnace atomic absorption spectrometry method. *Bull. Vet. Inst. Pulawy.*, 49: 89-92.
- Vanderpump, M and Tunbridge, M (2008). *Thyroid disease (The Facts)*. 2nd Edn., Oxford, Oxford University Press. P: 3.
- Yu, MH (2004). *Environmental toxicology: biological and health effects of pollutants*. 2nd Edn., Boca Raton, FL, CRC Press. P: 186.