



**IJVR**

ISSN: 1728-1997 (Print)  
ISSN: 2252-0589 (Online)

**Vol. 21**

**No.1**

**Ser. No.70**

**2020**

**IRANIAN  
JOURNAL  
OF  
VETERINARY  
RESEARCH**



## Short Paper

# Comparison of multiplex and ordinary PCR for diagnosis of paratuberculosis and tuberculosis in blood samples (buffy coat) of cattle and buffaloes

Brahma, D.<sup>1</sup>; Narang, D.<sup>2\*</sup>; Chandra, M.<sup>2</sup> and Singh, S. T.<sup>3</sup>

<sup>1</sup>MVSc Student in Veterinary Microbiology, Department of Veterinary Microbiology, College of Veterinary Science, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana-141004, Punjab, India; <sup>2</sup>Department of Veterinary Microbiology, College of Veterinary Science, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana-141004, Punjab, India; <sup>3</sup>Directorate of Livestock Farm, College of Veterinary Science, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana-141004, Punjab, India

\*Correspondence: D. Narang, Department of Veterinary Microbiology, College of Veterinary Science, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana-141004, Punjab, India. E-mail: [deeptivet@rediffmail.com](mailto:deeptivet@rediffmail.com)

(Received 4 Sept 2018; revised version 30 May 2019; accepted 24 Jun 2019)

## Abstract

**Background:** Paratuberculosis and tuberculosis (TB) caused by *Mycobacterium avium paratuberculosis* (MAP) and *Mycobacterium tuberculosis* complex (MTC), respectively are economically important, chronic debilitating diseases affecting the dairy herds and are also potential zoonotic threats. **Aims:** Differential diagnosis of paratuberculosis and TB in blood samples of cattle and buffaloes. **Methods:** In this study, an in-house developed multiplex polymerase chain reaction (PCR) targeting MAP, *Mycobacterium bovis* and *Mycobacterium smegmatis* was used in blood samples (buffy coat) parallel with IS900 PCR and *esxB* PCR for diagnosis of paratuberculosis and TB, respectively; in a total of 202 cattle and buffaloes. **Results:** Out of 202 animals, 12 (5.9%) and 17 (8.4%) animals were positive for MAP by multiplex PCR and IS900 PCR, respectively; from which only 8 (4%) animals were positive by both tests; whereas 4 and 9 animals were exclusively positive by multiplex PCR and IS900 PCR, respectively. None of the animals were found to be positive for *M. bovis* and *M. smegmatis* by the multiplex PCR. However, the *esxB* PCR detected 13 (6.4%) animals positive for TB. In fact, 3 (1.5%) animals were found to be co-infected by both paratuberculosis and TB. **Conclusion:** The in-house multiplex PCR detected MAP in buffy coat and there was a fair degree of agreement between the multiplex PCR and IS900 PCR in detection of MAP DNA though the latter detected more number of animals to be positive for MAP. Besides, *esxB* PCR showed a high diagnostic potential and can be used for diagnosis of TB from blood.

**Key words:** Blood samples, Multiplex PCR, Paratuberculosis, Tuberculosis

## Introduction

Paratuberculosis or Johne's disease (JD), caused by *Mycobacterium avium paratuberculosis* (MAP), is a chronic and progressive enteritis of domestic and wild ruminants, and is endemic in cattle of developing countries (OIE, 2019a). *Mycobacterium avium paratuberculosis* has impact on food safety and is also associated with Crohn's disease, which is a chronic inflammatory bowel disease in humans (Hermon-Taylor, 2009; Singh *et al.*, 2010; Carvalho *et al.*, 2012). Polymerase chain reaction (PCR) assays using primers specific for *F57*, *ISMav2*, *ISMAP02*, and *ISMAP04* elements have been used for specific detection of MAP DNA. However, the higher number of copies of IS900 element in comparison to other IS elements makes IS900-based detection very sensitive (Singh *et al.*, 2010; Carvalho *et al.*, 2012).

Tuberculosis (TB) is also an infectious chronic debilitating disease of zoonotic importance, affecting a wide range of livestock, domestic and wild animals, besides humans (OIE, 2019b). It is caused by the

pathogenic mycobacterial species, *Mycobacterium tuberculosis* complex (MTC) comprised of *Mycobacterium bovis*, *M. tuberculosis*, *Mycobacterium africanum*, *Mycobacterium canneti*, *Mycobacterium microti*, *Mycobacterium pinnipedii*, and *Mycobacterium caprae* (Olsen *et al.*, 2010). Tuberculosis in milch animal is mainly caused by *M. bovis* (OIE, 2009). Molecular diagnosis of TB using *esxB* (CFP-10) PCR targeting *esxB* gene present in the region of difference 1 (RD1) region of the pathogenic mycobacterial species, is a fast diagnostic tool having higher sensitivity and specificity (Dikshit *et al.*, 2012; Brahma *et al.*, 2017).

Opportunistic mycobacterial infections due to environmental mycobacteria may also occur in immunocompromised animals. One such example is the *Mycobacterium smegmatis*, which is a saprophytic, rapid growing, atypical, non-tuberculous mycobacteria (NTM) (Quinn *et al.*, 2011) and has the potential of causing opportunistic infection. *Mycobacterium smegmatis* has been reported from relapsing polygranulomatous mastitis in cattle (Siqueira *et al.*, 2016) and was also detected in milk samples among other atypical mycobacteria, which

may be a public health concern (Franco *et al.*, 2013; Bolaños *et al.*, 2017).

Therefore, compared to the traditional culture and microscopy method of diagnosis, the molecular diagnostic technique using Blood PCR is rapid and has a high potential for diagnosis of JD (Bhide *et al.*, 2006; Singh *et al.*, 2013), as well as TB (Zali *et al.*, 2014), especially during the bacteremic stage in subclinical or clinical stage. Therefore, in this study an in-house developed multiplex-PCR which differentiates mycobacterial infections caused by MAP, *M. bovis* and *M. smegmatis* was used in blood samples (buffy coat) of cattle and buffaloes for differential diagnosis of JD, TB and *M. smegmatis* infection. Further, this multiplex PCR was similarly complemented by conventional IS900 PCR and *esxB* (CFP-10) PCR for comparison of the diagnostic potential for MAP and MTC, respectively.

## Materials and Methods

### Collection of blood samples and DNA extraction

A total of 202 animals (42 cattle and 160 buffaloes) aged 2 years and above were selected randomly from an organized dairy farm in Ludhiana. Whole blood samples were collected and buffy coat was obtained from 1 ml each of whole blood samples by centrifugation at 12,000 g for 10 min and then washed with phosphate-buffered saline (PBS) for 2-3 times by re-centrifugation until white pellet is obtained. The buffy coat was subjected to DNA extraction using QIAamp DNA blood mini kit (Qiagen). The DNA sample was stored at -20°C until further use.

### In-house developed multiplex PCR

Multiplex PCR primers (Brahma *et al.*, 2017) designed with the help of *in-silico* PCR targeting three mycobacterial species: MAP, *M. bovis*, and *M. smegmatis* (Table 1) were used.

Brahma *et al.* (2017), assessed the sensitivity of the in-house multiplex primers by using ten-fold serial dilution of the standard DNA of *M. tuberculosis* (IMTECH, Chandigarh), MAP (GENEKAM, Germany) and *M. Smegmatis* (Microbiologics). Specificity of the

primers were cross-tested individually against genomic DNA of *M. tuberculosis*, MAP, *M. smegmatis*, and NTM species (*Mycobacterium fortuitum* and *Mycobacterium kansasii*) and non-mycobacterial species (*Brucella abortus*, *Pasteurella multocida* and *Escherichia coli*).

The DNA extracted from the buffy coat was amplified by in-house multiplex PCR primers and PCR conditions were followed as per Brahma *et al.* (2017).

### IS900 PCR

The DNA extracted from the buffy coat was also amplified by MAP species specific PCR based on the insertion sequence IS900. The sequences are as follows: Forward (IS900/150C): 5'- CCG CTA ATT GAG AGA TGC GAT TGG - 3' and Reverse (IS900/921): 5'- AAT CAA CTC CAG CAG CAG CGC GGC CTC G -3' designed to amplify a 229 bp target sequence (Vary *et al.*, 1990; Singh *et al.*, 2008). Reaction volume and PCR cycling conditions were followed as per Brahma *et al.* (2017).

### *esxB* (CFP-10) PCR

The primer sequences for CFP-10 were as follows: Forward: 5'-ATG GCA GAG ATG AAG ACC GAT GCC GCT-3' and Reverse: 5'-TCA GAA GCC CAT TTG CGA GGA CAG CGC C-3' (Dikshit *et al.*, 2012) giving a product size of 302 bp. Sample DNA from buffy coat was amplified by *esxB* (CFP-10) PCR, for detection of MTC. Reaction volume and PCR cycling conditions were followed as per Brahma *et al.* (2017) with a little modification, i.e., out of total 25 µL reaction volume, the DNA template was increased from 6 µL to 8 µL and nuclease free water was reduced from 4.5 µL to 2.5 µL, so as to increase the chances of detection, whereas the other reagent compositions were unchanged.

## Results

### In-house developed multiplex PCR


The sensitivity of *M. bovis*, MAP, and *M. smegmatis* primers were as little as 170 fg/µL, 300 fg/µL, and 51 fg/µL of genomic DNA, respectively. None of the organisms other than the specific standard DNA had

**Table 1:** Primer sequences for the in-house designed multiplex PCR

Target organism and strain		Primer sequence	Location of primer gene sequence	Size of PCR product
<i>M. avium</i> subsp. <i>paratuberculosis</i> MAP4 CP005928.1	Forward	5' - CGCGCGTACC TGACAAAAC - 3'	562055 - 562037	187 bp
	Reverse	5' - TCACCCTGAC ACTGACAGACA - 3'	561869 - 561889	
<i>M. bovis</i> strain SP38 CP015773.1	Forward	5' - GATGGTGGAA CACGACCACT - 3'	4138314 - 4138333	571 bp
	Reverse	5' - TTGATCGACC GTTCCGGTTT - 3'	4138865 - 4138884	
<i>M. smegmatis</i> MC2 155 CP009494.1	Forward	5' - ACCATGTCTAT CTCAGTGTGCT - 3'	3877883 - 3877904	628 bp
	Reverse	5' - ACGCTCGAGGT CCACTACAA - 3'	3878510 - 3878491	

amplification at the standardised annealing temperature i.e. 65.5-68°C which clearly indicates the specificity of in-house multiplex PCR (Brahma *et al.*, 2017).

Out of 202 animals, 12 (5.9%) were found to be positive for MAP infection by in-house multiplex PCR (Fig. 1). However, none of the animals were found to be positive for *M. bovis* and *M. smegmatis*.




**Fig. 1:** Amplification of DNA from the buffy coat using in-house developed multiplex primers. Lane M: 100 bp plus ladder. L1: Positive control (*M. avium paratuberculosis*, *M. tuberculosis*, and *M. smegmatis*), L2: Negative control, and L3-L14: Blood samples

### IS900 PCR

The analytical sensitivity of the IS900 PCR was upto 30 fg/ $\mu$ L of MAP DNA (GENEKAM, Germany) and no other bacteria other than the target species was detected indicating its specificity (Brahma *et al.*, 2017).

Out of a total 202 animals screened for JD, a total of 17 (8.4%) animals were positive by IS900 PCR (Fig. 2). However, only 8 (4%) out of 17 animals were positive by both the multiplex and IS900 PCR; whereas 4 and 9 animals were exclusively positive by multiplex PCR and IS900 PCR, respectively.




**Fig. 2:** Amplification of DNA from the buffy coat using IS900 primers. Lane M: 100 bp plus ladder. L1: Positive control (*M. avium paratuberculosis*), L2: Negative control, and L3-L10: Blood samples

### esxB (CFP-10) PCR

The analytical sensitivity of *esxB* (CFP-10) primers were upto 800 fg/ $\mu$ L of *M. tuberculosis* DNA (IMTECH, Chandigarh) and none of the NTM or other bacteria were detected (Brahma *et al.*, 2017).

Out of the 202 animals screened for TB, only 13 (6.4%) animals were found to be positive for TB by *esxB* PCR (Fig. 3). In fact, the combined results of IS900 and *esxB* PCR revealed incidences of co-infection with both JD and TB in 3 animals which had debilitated health condition.



**Fig. 3:** Amplification of DNA from the buffy coat using *esxB* (CFP-10) primers. Lane M: 100 bp plus ladder. L1-L4: Blood samples, L5: Positive control (*M. tuberculosis*), and L6: Negative control

### Discussion

In this study, the in-house developed multiplex PCR detected MAP in blood samples though it did not detect *M. bovis* and *M. smegmatis*. In contrast, Brahma *et al.* (2017), using the same multiplex PCR detected both MAP and *M. bovis* from tissue samples, though the sensitivity for detection of *M. bovis* was comparatively less than the *esxB* PCR. Amplification of *M. bovis* in multiplex PCR may be inhibited in blood samples due to PCR inhibitors or it could be presence of DNA in undetectably low concentration or due to presence of low copy number of the target sequence. In other studies by different workers, the accuracy of multiplex PCR for differential detection of *M. bovis* and *M. tuberculosis* was found to be 100% in terms of specificity and could detect as little as 20 pg of genomic DNA (Shah *et al.*, 2002; Bakshi *et al.*, 2005).

IS900 PCR-based detection of MAP from white blood cells (WBCs) has also been reported by some workers (Naser *et al.*, 2004; Stott *et al.*, 2005). The reason for detection of MAP in more number of samples by IS900 PCR compared to the in-house developed multiplex PCR could be due to the higher copy number of IS900 element as reported by Vary *et al.* (1990) and Singh *et al.* (2010). However, using tissue samples, Brahma *et al.* (2017) detected MAP DNA in 2 out of 2 histopathologically JD positive cases by the in-house multiplex PCR.

In other studies, it has been found that MAP could be readily detected in blood from cattle with disseminated infections, and possibly in the blood of infected young and asymptomatic animals (Juste *et al.*, 2005). Blood PCR targeting IS900 gene was rapid, highly sensitive, and specific for detecting MAP infection at any stage of infection in any age of goats (Singh *et al.*, 2010).

*Mycobacterium avium paratuberculosis* DNA was also detected in WBCs from about 11% of 262 Indian cattle of unspecified age or disease status (Bhide *et al.*, 2006). Singh *et al.* (2013) found substantial agreement between “microscopy” and blood PCR detecting 39.3 and 13.1% positive by faecal microscopy and blood PCR, respectively; though Blood PCR detected more cases from heavy MAP shedders in microscopy. It may be assumed that low positivity in blood PCR may be due to presence of MAP in blood stream for a limited period (Singh *et al.*, 2013). Gwozdz *et al.* (2000) contrarily showed poor performance of “blood PCR” to detect subclinically infected sheep. Barrington *et al.* (2003) had also recorded lower sensitivity of “blood-PCR” in comparison to PCR applied on milk, liver and faecal samples of advanced subclinically infected cows. The overall sensitivity of blood PCR for JD diagnosis from different studies was 11-66%. However, Youssef *et al.* (2014), reported that, out of 150 faecal samples, 31.33%, 29.33%, and 42.66% were found positive whereas MAP DNA was detected in only 4.67%, 3.33%, and 5.33% of milk samples by conventional IS900 PCR, conventional F57 PCR and IS900 real-time (RT) PCR, respectively. Gümüşsoy *et al.* (2015), in PCR of milk and stool samples, detected MAP DNA in 20 (13.61%) and 42 (28.57%) samples, respectively. However, MAP is only shed via milk in a small proportion of cows with subclinical JD for a limited period of time (Khol *et al.*, 2013) and the shedding of MAP in faeces and milk is not synchronized (Gao *et al.*, 2009).

Several PCR systems have also been developed for the diagnosis of TB, e.g., PCR amplification of *esxA* and *esxB* genes targeting ESAT-6 and CFP-10 proteins, respectively present in pathogenic mycobacterial species can be used for detection of *M. tuberculosis* as well as *M. bovis* (Pinxteren *et al.*, 2000; Rogerson *et al.*, 2006; Dikshit *et al.*, 2012). Using *esxB* PCR on tissue samples Brahma *et al.* (2017) detected all 7 out of 7 histopathologically TB positive cases, indicating its efficiency in TB diagnosis. Zali *et al.* (2014), using JB21 and JB22 primers specific for *M. bovis*, detected *M. bovis* positive in 8% of blood and lymph node samples of cattle. In this study, PCR targeting *esxB* (CFP-10) gene in blood samples detected 13 (6.4%) TB positive animals which were not detected by the in-house multiplex PCR, thus, indicating the early diagnostic potential of *esxB* (CFP-10) PCR. After all, 3 of the TB infected animals found to be positive for JD indicates presence of co-infection with both the diseases within the herd. Infection of either JD or TB may make the animals more susceptible to co-infection due to the immunocompromised condition (Brahma *et al.*, 2017).

In conclusion, “Blood PCR” using buffy coat was rapid, highly sensitive and specific, and the combination of various PCR techniques increased the chances of successful diagnosis of both JD and TB infection in cattle and buffaloes. The use of multiplex-PCR was able to detect MAP and differentiate from other mycobacterial infections in blood samples, thus providing a rapid diagnosis and its confirmation for the

specific disease of concern, JD. However, the probable limitation is that, a large-scale study is required to determine whether this PCR assay is adequate for JD and TB control program.

## Acknowledgements

The authors are very grateful to DBT (Department of Biotechnology), Government of India for providing funds for the present work through a Project (BT/PR5776/MED/30/928/2012) and to the Director of Research, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana for providing the necessary facilities.

## References

- Bakshi, CS; Shah, DH; Verma, R; Singh, RK and Malik, M (2005). Rapid differentiation of *Mycobacterium bovis* and *Mycobacterium tuberculosis* based on a 12.7-kb fragment by a single tube multiplex-PCR. *Vet. Microbiol.*, 109: 211-216.
- Barrington, GM; Gay, JM; Eriks, IS; Davis, WC; Evermann, JF; Emerson, C; O'Rourke, JL; Hamilton, MJ and Bradway, DS (2003). Temporal patterns of diagnostic results in serial samples from cattle with advanced paratuberculosis infections. *J. Vet. Diagn. Invest.*, 15: 195-200.
- Bhide, M; Chakurkar, E; Tkacikova, L; Barbudhe, S; Novak, M and Mikula, I (2006). IS900-PCR-based detection and characterization of *Mycobacterium avium* subsp. *paratuberculosis* from buffy coat of cattle and sheep. *Vet. Microbiol.*, 112: 33-41.
- Bolaños, CAD; de Paula, CL; Guerra, ST; Franco, MMJ and Ribeiro, MG (2017). Diagnosis of mycobacteria in bovine milk: an overview. *Rev. Inst. Med. Trop. São Paulo.* 59: e40.
- Brahma, D; Narang, D; Chandra, M; Gupta, K; Singh, A and Kaur, G (2017). Diagnosis of Mycobacterial infections (Tuberculosis and Paratuberculosis) in tissue samples using molecular (inhouse multiplex PCR, PCR and TaqMan real-time PCR), histopathology and immunohistochemical techniques. *Trop. Biomed.*, 34: 1-17.
- Carvalho, IA; Vinicius, CEB; Souza, IM; Zanardo, LG; Filho, JDR; Gomes, MJP and Moreira, MAS (2012). Diagnosis of Paratuberculosis in cattle: microbiological culture, serology and PCR. *Braz. J. Microbiol.*, 43: 581-585.
- Dikshit, M; Sharma, RJ; Adsool, AD and Chaphalkar, SR (2012). ESAT-6 and CFP-10 proteins of *Mycobacterium tuberculosis* in making diagnostic tool for TB. *J. Biotechnol. Lett.*, 3: 28-30.
- Franco, MM; Paes, AC; Ribeiro, MG; Pantoja, JC; Santos, AC; Miyata, M; Leite, CQ; Motta, RG and Listoni, FJ (2013). Occurrence of mycobacteria in bovine milk samples from both individual and collective bulk tanks at farms and informal markets in the southeast region of Sao Paulo, Brazil. *BMC Vet. Res.*, 9: 85.
- Gao, A; Odumeru, J; Raymond, M; Hendrick, S; Duffield, T and Mutharia, L (2009). Comparison of milk culture, direct and nested polymerase chain reaction (PCR) with fecal culture based on samples from dairy herds infected with *Mycobacterium avium* subsp. *paratuberculosis*. *Can.*

- J. Vet. Res., 73: 58-64.
- Gümüşsoy, KS; İça, T; Abay, S; Aydın, F and Hizlisoy, H** (2015). Serological and molecular diagnosis of paratuberculosis in dairy cattle. *Turk. J. Vet. Anim. Sci.*, 39: 147-153.
- Gwozdz, JM; Thompson, KG; Manktelow, BW; Murray, A and West, DM** (2000). Vaccination against paratuberculosis of lambs already infected experimentally with *Mycobacterium avium* subspecies *paratuberculosis*. *Aust. Vet. J.*, 78: 560-566.
- Hermon-Taylor, J** (2009). *Mycobacterium avium* subspecies *paratuberculosis*, Crohn's disease and the Doomsday scenario. *Gut Pathog.*, 1: 15. doi: 10.1186/1757-4749-1-15.
- Juste, RA; Garrido, JM; Geijo, MV; Elguezabal, N; Aduriz, G; Atxaerandio, R and Sevilla, I** (2005). Comparison of blood PCR and ELISA for detection of *Mycobacterium avium* subsp. *paratuberculosis* infection in sheep and cattle. *J. Vet. Diagn. Invest.*, 17: 354-359.
- Khol, JL; Wassertheurer, M; Sodoma, E; Revilla-Fernández, S; Damoser, J; Österreicher, E; Dünser, M; Kleb, U and Baumgartner, W** (2013). Long-term detection of *Mycobacterium avium* subspecies *paratuberculosis* in individual and bulk tank milk from a dairy herd with a low prevalence of Johne's disease. *J. Dairy Sci.*, 96: 3517-3524.
- Naser, SA; Ghobrial, G; Romero, C and Valentine, JF** (2004). Culture of *Mycobacterium avium* subspecies *paratuberculosis* from the blood of patients with Crohn's disease. *Lancet*. 364: 1039-1044.
- OIE** (2009). Chap. 2.4.6: Bovine tuberculosis. *Terrestrial Manual*. World Organization for Animal Health, Paris, France.
- OIE** (2019a). <http://www.oie.int/en/animal-health-in-the-world/animal-diseases/Paratuberculosis/> Accessed on 04.02.2019.
- OIE** (2019b). <http://www.oie.int/en/animal-health-in-the-world/animal-diseases/Bovine-tuberculosis/> Accessed on 04.02.2019.
- Olsen, I; Barletta, RG and Thoen, CO** (2010). *Mycobacterium*. In: Gyles, CL; Prescott, JF; Songer, JG and Thoen, CO (Eds.), *Pathogenesis of bacterial infections in animals*. (4th Edn.), Ames, Iowa, Wiley-Blackwell. PP: 113-132.
- Pinxteren, LAHV; Ravn, P; Agger, EM; Pollock, J and Andersen, P** (2000). Diagnosis of tuberculosis based on the two specific antigens ESAT-6 and CFP-10. *Clin. Diagn. Lab. Immunol.*, 7: 155-160.
- Quinn, PJ; Markey, BK; Leonard, FC; Fitzpatrick, ES; Fanning, S and Hartigan, PJ** (2011). *Mycobacterium* species. In: Quinn, PJ; Markey, BK; Leonard, FC; Fitzpatrick, ES; Fanning, S and Hartigan, PJ (Eds.), *Veterinary microbiology and microbial disease*. (2nd Edn.), Chapter 23, Oxford, Wiley-Blackwell. PP: 250-262.
- Rogerson, BJ; Jung, YJ; LaCourse, R; Ryan, L; Enright, N and North, RJ** (2006). Expression levels of *Mycobacterium tuberculosis* antigen encoding genes versus production levels of antigen specific T cells during stationary level lung infection in mice. *Immunology*. 118: 195-201.
- Shah, DH; Verma, R; Bakshi, CS and Singh, RK** (2002). A multiplex-PCR for the differentiation of *Mycobacterium bovis* and *Mycobacterium tuberculosis*. *FEMS Microbiol. Lett.*, 214: 39-43.
- Singh, SV; Singh, PK; Gupta, S; Chaubey, KK; Singh, B; Kumar, A; Singh, AV and Kumar, N** (2013). Comparison of microscopy and blood-PCR for the diagnosis of clinical Johne's disease in domestic ruminants. *Iran. J. Vet. Res.*, 14: 345-349.
- Singh, PK; Singh, SV; Kumar, H; Sohal, JS and Singh, AV** (2010). Diagnostic application of IS900 PCR using blood as a source sample for the detection of *Mycobacterium avium* subspecies *paratuberculosis* in early and subclinical cases of caprine paratuberculosis. *Vet. Med. Int.*, Article ID 748621, 8 pages.
- Singh, SV; Singh, AV; Singh, R; Sharma, S; Shukla, N; Mishra, S; Singh, PK; Sohal, JS; Kumar, H; Patil, PK; Misra, P and Sandhu, KS** (2008). Sero-prevalence of bovine Johne's disease in buffaloes and cattle population of north India using indigenous ELISA kit based on native *Mycobacterium avium* subsp. *paratuberculosis* 'Bison type' genotype of goat origin. *Comp. Immunol. Microbiol. Infect. Dis.*, 31: 419-433.
- Siqueira, FM; Lopes, CE; Snell, GG and Gomes, MJP** (2016). Identification of *Mycobacterium smegmatis* in bovine Mastitis. *Acta Sci. Vet.*, (Suppl. 1), 44: 166.
- Stott, AW; Jones, GM; Humphry, RW and Gunn, GJ** (2005). Financial incentive to control paratuberculosis (Johne's disease) on dairy farms in the United Kingdom. *Vet. Rec.*, 156: 825-831.
- Vary, PH; Anderson, PR; Green, E; Taylor, JH and McFadden, JJ** (1990). Use of highly specific DNA probes and the polymerase chain reaction to detect *Mycobacterium paratuberculosis* in Johne's disease. *J. Clin. Microbiol.*, 28: 933-937.
- Youssef, DGS; Sallam, FA; Darwish, SF and Amin, AS** (2014). Evaluation of conventional and real-time PCR assays for molecular diagnosis of Johne's disease in dairy cattle. *Int. J. Curr. Microbiol. App. Sci.*, 3: 969-981.
- Zali, MHS; Farajnia, F; Yahyapour, H; Moslemzadeh, T and Hashempour, A** (2014). Detection of *Mycobacterium bovis* in cattle suspected to tuberculosis by PCR method in Urmia-Iran. *BEPLS.*, 3: 152-157.